

# **THE SITE-SPECIFIC EFFECTS OF KINDLING ON COGNITION AND ADULT HIPPOCAMPAL NEUROGENESIS**

A Thesis submitted to the College of Graduate and Postdoctoral studies  
in Partial Fulfillment of the Requirements for the Degree of Master of Science in the Department  
of Medicine at the University of Saskatchewan

By Nikita Nogovitsyn

© Copyright Nikita Nogovitsyn, January 2017. All rights reserved.

## **PERMISSION TO USE**

In presenting this thesis/dissertation in partial fulfillment of the requirements for a Postgraduate degree from the University of Saskatchewan, I agree that the Libraries of this University may make it freely available for inspection. I further agree that permission for copying of this thesis/dissertation in any manner, in whole or in part, for scholarly purposes may be granted by the professor or professors who supervised my thesis/dissertation work or, in their absence, by the Head of the Department or the Dean of the College in which my thesis work was done. It is understood that any copying or publication or use of this thesis/dissertation or parts thereof for financial gain shall not be allowed without my written permission. It is also understood that due recognition shall be given to me and to the University of Saskatchewan in any scholarly use which may be made of any material in my thesis/dissertation.

Requests for permission to copy or to make other uses of materials in this thesis/dissertation in whole or part should be addressed to:

Head of the Department of Medicine

University of Saskatchewan

Saskatoon, Saskatchewan S7N5E5

Canada

«До тех пор, пока вы предмет не постигли, он для вас представляется сложным и туманным. Но как только истина уловлена, все становится простым. Признак истины - простота, и все гении просты своими истинами.»

Иван Петрович Павлов

(Лекция 1918)

“As long as you don’t understand the subject, the picture seems to be cloudy and perplexing. But once the truth is caught – everything becomes simple. Simplicity is the ultimate sign of the truth, and all geniuses are simple in their truths.”

Ivan Pavlov

(Lecture notes 1918)

## ABSTRACT

Nearly 1.5% of the general population is affected by epilepsy. Despite a long history of research and clinical endeavors to combat the disease, in 1 of 3 cases, epilepsy is intractable and resistant to medication treatment. Current pharmacological strategies target seizure onset as the primary manifestation of illness; however, patients with advanced stages of the disease suffer from a wide array of psychiatric comorbidities. Anxiety, depression and memory deterioration are the chief complaints of epileptic patients delegated from neurologic to psychiatric care.

Temporal lobe epilepsy is the most resistant form of epilepsy and the form that is most complicated by the presence of psychiatric and cognitive comorbidities that are increasingly recognized as critical factors in long-term patient care. These comorbidities are independent risk factors for poor quality of life. In fact, studies have shown that in patients with epilepsy, comorbid factors correlate more strongly with poor quality of life than does seizure frequency. At present, the behavioural comorbidities associated with temporal lobe epilepsy are poorly understood, and their management is difficult because many commonly prescribed anticonvulsant drugs make them worse. Therefore, even patients with some degree of seizure control often continue to experience debilitating behavioural problems that impair their daily living.

**This thesis attempts to understand the impact of seizures originating from different brain sites.** The research described in this thesis was conducted using an animal model called kindling to investigate the hypothesis that seizure-induced alterations in hippocampal neurogenesis play a significant role in the cognitive deficits associated with epilepsy. The experiment used 25 adult male rats. Rats were divided into four groups: hippocampal kindled (n = 6), amygdala kindled (n = 7), caudate nucleus kindled (n = 6) and control rats (n = 6). Kindled

rats received 99 electrical stimulations delivered to the appropriate brain region through an implanted bilateral electrode. At the end of the kindling phase, rats were subjected to a fear conditioning paradigm to assess cognitive behavior. After the fear conditioning, rats were sacrificed and immunohistochemical analyses were done to assess the impact of seizures on hippocampal neurogenesis and neuronal activation. The results indicated that there is a significant deterioration of cognitive performance following long-term limbic kindling (i.e., hippocampal and amygdala kindling only). This was accompanied by an increase in hippocampal neurogenesis that was paralleled with low expression of immediate early genes. Collectively, these findings enhance our understanding of mechanisms underlying the behavioral comorbidities associated with temporal lobe epilepsy and **demonstrate the link between adult neurogenesis and cognitive impairments following long-term kindling of limbic brain regions.**

## AKNOLEDGEMENTS

First, I would like to address my appreciation and gratitude to my research supervisor, Dr. Lisa Kalynchuk.

Dr. Kalynchuk, thank you for opening the door into the beautiful world of science. I am grateful for the opportunity to join your research team. You made a significant impact on my career path and provided me with the invaluable opportunity to develop on multiple levels. I highly appreciate your supervision, mentorship and guidance within my Masters program. I hold the deepest appreciation for your belief in my potential as a scientist. It has been a privilege to work and study in your laboratory.

I would also like to express many thanks to Dr. Caruncho for his valuable research advice. It was always a pleasure to talk to Dr. Caruncho about science and receive his optimistic attitude toward research endeavors.

Many thanks to Dr. Botterill for his time and patience teaching me and training me in the laboratory techniques that allowed me to successfully collect and analyze the data for my thesis.

I would also like to say thank you to my dear friend and lab mate, Milann Mitchell, who was an indispensable satellite within my graduate program; thank you for your support and friendship.

I would also like to thank my dear friend, Dr. Maryana Kravtsenyk. Thank you for being a paragon of inspiration, electrifying perseverance and unshakable faith in success.

Many thanks to all the members of my committee.

## TABLE OF CONTENTS

<b>PERMISSION TO USE.....</b>	<b>I</b>
<b>ABSTRACT.....</b>	<b>III</b>
<b>AKNOLEDGEMENTS .....</b>	<b>V</b>
<b>LIST OF ABBREVIATIONS .....</b>	<b>IX</b>
<b>LIST OF FIGURES .....</b>	<b>X</b>
<b>CHAPTER 1 .....</b>	<b>1</b>
<b>1.1 Epilepsy .....</b>	<b>1</b>
1.1.1. Epilepsy comorbidities.....	4
<b>1.2. Kindling .....</b>	<b>5</b>
<b>1.3. Neurogenesis .....</b>	<b>8</b>
<b>1.4 Neurogenesis and pathology .....</b>	<b>13</b>
<b>1.5. Markers of neurogenesis and cell activity .....</b>	<b>15</b>
<b>1.6 Neurogenesis markers .....</b>	<b>16</b>
1.6.1. FOS .....	17
1.6.2. ARC .....	18
<b>1.7. Neuroanatomy .....</b>	<b>19</b>
1.7.1. Hippocampus .....	19
1.7.2. Basolateral Amygdala.....	22
1.7.3. Caudate Nucleus .....	24
<b>1.8. Pavlovian Fear Trace Conditioning.....</b>	<b>25</b>
<b>1.9. Experimental goal and the hypothesis.....</b>	<b>27</b>
1.9.1. The specific aims and goals .....	27
<b>CHAPTER 2 .....</b>	<b>29</b>
<b>2.1. Abstract .....</b>	<b>30</b>
<b>2.2. Introduction .....</b>	<b>31</b>
<b>2.3. Materials and methods.....</b>	<b>33</b>
2.3.1. Subjects .....	33
2.3.2. Surgery .....	33
2.3.3. Kindling .....	36
2.3.4. BrdU injections .....	37
2.3.5. Fear conditioning .....	37
2.3.6. Perfusions.....	39
2.3.7. Post-mortem analysis: immunohistochemistry procedures.....	39
2.3.7.1. BrdU staining .....	39
2.3.7.2. Fos staining .....	40
2.3.7.3. Arc staining.....	41
2.3.7.4. Immunofluorescence double labelling .....	41

2.3.7.5. Doublecortin immunohistochemistry.....	42
2.3.7.6. Quantification.....	43
2.3.7.7. Statistical analyses .....	45
<b>2.4. Results.....</b>	<b>46</b>
2.4.1. Kindling impacts fear trace conditioning.....	46
2.4.2. Stereology counting for BrdU as a marker of neurogenesis .....	49
2.4.3. Doublecortin fluorescence .....	51
2.4.4. Profile counting for Arc immunoreactivity.....	53
2.4.5. Profile counting for Fos immunoreactivity .....	55
2.4.6. Double labelling for BrdU and Arc.....	58
2.5. Discussion .....	62
<b>CHAPTER 3.....</b>	<b>68</b>
<b>3.1. General discussion.....</b>	<b>68</b>
3.1.1. Aberrant neurogenesis and cognition.....	69
3.1.2. Why do seizures impede neuronal activation?.....	70
<b>3.2. Limitations .....</b>	<b>72</b>
<b>3.3. Future directions .....</b>	<b>75</b>
<b>REFERENCES.....</b>	<b>78</b>

## LIST OF ABBREVIATIONS

dHip – dorsal hippocampus  
BLA – basolateral amygdala  
BrdU – bromodeoxyuridine  
CS – conditioned stimulus  
TBS – tris-buffered saline  
PBS – phosphate-buffered saline  
IHC – immunohistochemistry  
ARC – activity-regulated cytoskeleton-associated protein, also known as Arg3.  
DCX – doublecortin  
NHS – normal horse serum  
NGS – normal goat serum  
BSA – bovine serum albumin  
GCL – granule cell layer  
DG – dentate gyrus  
OB – olfactory bulb  
AN – adult neurogenesis  
FTC – fear trace conditioning  
IEG – immediate early gene  
TLE – temporal lobe epilepsy  
BDNF – brain neurotrophic factor  
CN – caudate nucleus  
DS – dorsal striatum  
SN – substantia nigra (lat.)  
CeA – central amygdala

## LIST OF FIGURES

**Figure 1.** Representative photomicrograph of Nissl stained brain tissue

**Figure 2.** Kindling affects fear trace conditioning

**Figure 3.** Comparative photomicrograph of BrdU expression across treatment groups

**Figure 4.** Quantification analysis of BrdU expression in SGZ and GCL zones (stereology)

**Figure 5.** Illustrative example of DCX immunofluorescence across treatment groups

**Figure 6.** Comparative photomicrograph of Arc expression in the dentate gyrus

**Figure 7.** Effect of kindling on Arc expression following context fear memory retrieval

**Figure 8.** Comparative illustration of FOS expression across across treatment groups

**Figure 9.** Kindling impacts FOS expression in limbic-kindled rats

**Figure 10.** Pattern of fluorescent immunoreactivity (BrdU+Arc) across treatment groups

**Figure 11.** Illustrative example of colocalized cell. Merged images of BrdU and Arc

**Figure 12.** The effect of limbic kindling on BrdU and Arc fluorescent immunostaining

**Figure 13.** Colocalization analysis of merged images in immunofluorescent double labeling

# CHAPTER 1

## GENERAL INTRODUCTION

### 1.1 Epilepsy

Epilepsy is a chronic neurological disease affecting approximately 50 million people worldwide (Birbeck et., 2010). It is also estimated that an additional 2.4 million people will be diagnosed with epilepsy every year. The majority of diagnoses (80%) fall onto patients from developing countries and importantly, it affects age groups during their most productive years of life (Birbeck et al., 2010). One in ten people would experience at least one epileptic seizure within the normal lifespan and one third of them will develop epilepsy.

The current International Statistical Classification of Diseases (ICD – 10) provides following definition of epilepsy:

*‘Epilepsy is a brain disorder characterized by episodes of abnormally increased neuronal discharge resulting in transient episodes of sensory or motor neurological dysfunction, or psychic dysfunction (ICD – 10: G-40).’* ICD – 10 dedicates nearly 70 diagnostic chapters under the umbrella of ‘epilepsy’. This is a heterogeneous disease complicated by a wide variety of clinical manifestations that depend upon many factors such as brain site, etiology, duration and time pattern of the onset. Such wide array of factors may have synergetic effects and complicate the process of diagnosis and treatment.

There are numerous causes of epilepsy, as both genetic and acquired factors may contribute and interact, suggesting that the pathophysiology of this disease is quite complex. Spontaneous seizures, which are the definitive feature of epilepsy leading to a diagnosis, are thought to be the result of a disbalance between inhibition and excitation shifting the system towards hyperexcitability. This process may occur in a restricted brain site and manifest as a

partial epilepsy, and/or subsequently ramify into a generalized epilepsy when seizure activity synchronizes and hyperpolarizes several brain sites simultaneously. Genetic causes are thought to be the most widespread causative effect for epileptic seizure development. Brain tumors, either benign or malignant are another common cause for epilepsy, accounting for nearly 30% of epilepsy cases (Liigant et al., 2001). Brain injury is another risk factor that may unfold into epilepsy in 6% of cases (Hausser et al., 1993) and generalized seizures are associated with traumatic brain injuries within the onset of the seizure (Lowenstein et al., 2009). Stroke is a major risk factor for elderly patients and may explain epilepsy occurrence in one-third of patients (Cleary et al., 2004). The list of epilepsy etiology may be associated with infections like toxoplasmosis or meningitis, febrile seizures in newborns or those seizures occurring in obstetric patients under the condition of severe eclampsia. Despite such variability, there appears to be a common pathophysiological mechanism. All epileptic seizures arise from excessively synchronous activity and sustained discharge of a selected neuronal population (Fisher et al., 2005). Therefore, the unifying feature of all epileptic syndromes is a persistent increase of excitatory systems. Albeit, because we do not have a full and clear picture of the mechanisms that underlie this excitation, further scientific investigation is necessary.

As mentioned above, spontaneous and unprovoked seizure onset is the definitive diagnostic feature of epilepsy. These seizures may be presented as brief episodes of a loss of consciousness accompanied by involuntary movement involving a part of the body (partial) or the entire body (generalized). Patients suffering from epilepsy also demonstrate increased risk for other psychiatric disorders that are paralleled with cognitive disturbances. Pharmacological resistance is a common phenomenon among patients with full-blown epilepsy converting them into candidates for surgery. However surgical resection of affected brain sites does not alleviate

cognitive deficits. Further, surgical patients often have more than one brain lesion within the affected area, a phenomenon called ‘dual pathology’. This appears to be common in younger patients. Additionally, current pharmacological treatment has a variety of side effects that contribute towards cognitive deterioration in patients with epilepsy. Alas, **nowadays treatment strategies leave patients with no effective treatment against the cognitive and emotional comorbidities of epilepsy.**

Temporal lobe epilepsy (TLE) is the most common form of epilepsy in adults. It is also one of the most pharmaco-resistant types and appears to be hard to access for surgical treatment. The diagnosis of TLE may be further divided into partial with or without consciousness impairments and complex partial subtypes. Detection of epileptic foci is an additional diagnostic step that may label TLE with limbic or neocortical subtype, which is a rare clinical case scenario. Low amplitude with rapid activity rhythmic spikes, either synchronous or asynchronous, in the electroencephalogram (EEG) is a defining feature of this diagnostic group. The specific anatomic position of the hippocampus puts it into a crucial role contributing towards the progression of limbic TLE. Post-mortem analysis of patients suffering from TLE indicates total cellular reorganization in the hippocampal formation as sclerotic changes take place (Parent et al., 1997). Besides possible consciousness impairments, seizures can be accompanied by a variety of sensory phenomena such as auras, hallucinations or persistent Deja vu feelings. Epileptic onset frequently involves repetitive behavioural patterns or orofacial automatisms (head nodding, tremor of the limbs, lip smacking). The seizure itself is a relatively short event that leaves the patient in a post-ictal period. Interestingly, to date, there is no clear understanding about how and why seizures are terminated. However, the onset itself is just a visible manifestation of

progressive changes happening within the whole system. The chronic progression of behavioural changes in epileptic patients leads to a lower quality of life and stigmatization.

### **1.1.1. Epilepsy comorbidities**

A comorbid condition is one that occurs during the progression of a main disease such as epilepsy. Thus, psychiatric and social comorbidities are diagnostic satellites of epilepsy as an index disease. Nearly 50% of patients suffering from epilepsy report at least one psychiatric or somatic comorbid disorder (Keezer et al., 2015; Forsgren et al., 1992). Among them, 20% of patients suffering from TLE demonstrate psychiatric complaints (Gaitatzis et al., 2004). Mood disorders, anxiety, psychosis and personality changes are the most common conditions in patients with epilepsy (Bragatti et al., 2004). Importantly, depression has a much more debilitating effect on the quality of life in patients with TLE than seizure frequency itself (Boylan et al., 2004; Perrine et al., 1995). Subsequently depressive states are the strongest risk factors for suicide in patients with epilepsy (Schmitz et al., 2005). Fear and anxiety are also common complaints as they are the most frequent feelings induced by seizures. In fact, rapid onset of fear and anxiety may present as panic disorder, leading to misdiagnosis (Sazgar et al., 2003).

Patients with TLE also suffer from cognitive impairments because the medial temporal lobes are involved in learning and memory processes. Patients with TLE often demonstrate deficits in declarative and episodic memory (Butterbaugh et al., 2004; Schwarcz et al., 2002), as well as impairments in long-term memory consolidation and spatial memory (Cataldi et al., 2013; Haag et al., 2010).

Any comorbid state can affect therapeutic decisions in people with epilepsy. Migraines, hepatic insufficiency or depression could be factors influencing the decision toward a type of antiepileptic drug as a treatment. Cognitive deficits may also question the suitability of surgical intervention in patients with TLE because of potential severe memory impairments from temporal resections.

Taking into consideration the above discussion, epilepsy may present not only as a neurological disease but also as a psychiatric condition. The clinician may be deceived by depressive or affective symptoms that mask epilepsy, thus complicating diagnostic procedures. The patient, on the other side, may face therapeutic ‘Zugzwang’ in attempt to combat epileptic comorbidity because neither pharmacological nor surgical intervention are able to reduce psychiatric impairments. Future scientific efforts are needed to understand the underlying mechanisms that govern epilepsy progression. It is evident and clear that current progress in the treatment of epileptic patients is not satisfactory, and specific need exists to better understand the mechanisms underlying behavioral co-morbidities in epilepsy and potentially bring novel treatment strategies to the clinic.

## **1.2. Kindling**

In 1969, Graham V Goddard and colleagues released a publication titled “A permanent change in brain function resulting from daily electrical stimulation” where they provided a detailed description of a possible animal model of epilepsy. ‘Kindling’ is a term coined by those pioneering researchers and it refers to the gradual development and intensification of motor seizures resulting from daily electrical stimulation of a specific brain region. To target a selected brain site, animals (typically rats) are first subjected to stereotaxic surgery aiming to implant

permanent electrodes to deliver electrical stimulations. Intensification of seizure activity that is paralleled with behavioural convulsions resulting from electrical stimulation is a reliable hallmark of a successful kindling procedure. The kindling phenomenon proved to be a valid animal model of epilepsy once it was demonstrated that long-term electrical stimulation may elicit spontaneous convulsions (Pinel et al., 1978). A variety of anticonvulsant drugs may decrease and even reverse epileptic activity in limbic-kindled rats (Adamec et al., 1986) – that was another property of kindling that confirmed its relevance to kindling as a model of epilepsy. Much like current epileptogenic dogma, kindling decreases protective brain mechanisms (i.e., the GABA system) in response to focal discharge (Kokaia et al., 1994). This brief description clearly demonstrates that kindling initially models focal partial seizures. With the daily progression of electrical stimulation and the recruitment of other brain regions into seizure discharges, kindling comes to model complex partial seizures with secondary generalization (McIntyre et al., 1979). Finally, animals become undeniably epileptic once seizures become spontaneous after many days of electrical stimulations (Pinel et al., 1975; Wada et al., 1976).

Kindling creates a “pathology” on multiple levels. Goddard, as pioneer of kindling, argued that kindling induces structural alterations within or between neurons (Goddard et al., 1969). However, nowadays kindling is thought to systematically induce a sequence of long-term functional and structural changes, initiated by perturbations in synaptic transmission that eventually lead to morphological changes within neurons and neuronal circuits (Sutula et al., 1991). The initial episode of network synchronization during kindling upregulate NMDA receptor dependent components (Sayin et al., 1989). Then, NMDA receptor dependent elements induce intracellular  $Ca^{2+}$  influx. A high intracellular  $Ca^{2+}$  state activates the transcription of early immediate genes, which results in an avalanche of gene alterations (Hughes et al., 1998; Shin et

al., 1990; Sutula et al., 1996). Kindling also affects neurotrophins such as BDNF, NT3, NT4 (Xiao-Ping et al., 2004). In sum, these 'subneuronal' alterations contribute towards cellular pathology, and finally neuronal cell loss and apoptosis emerge.

Even a single seizure induced by kindling can induce apoptosis (Bengzon et al., 1997). Systematically repeated electric stimulations result in a measureable neuronal loss (Kavazos et al., 1994; Kolotski et al., 2002). Neuronal cell death is followed by a compensatory mechanism of 'tissue replacement', thus glial cells replace previously functional neurons. And finally, prolonged and systematic cellular death results in a pathomorphological phenomenon called hippocampal sclerosis (Thom et al., 2014). Therefore, seizure-induced neuronal injury and subsequent neuronal loss with hippocampal sclerosis initially starts from excitotoxicity, excessive Na<sup>+</sup> and Ca<sup>2+</sup>, osmolytic stress and free radical production (Henshall et al., 2007).

Once intrinsic compensatory mechanisms approach exhaustion, functional dysregulation can occur. This can be seen in kindling as a change in multiple domains of behaviour. For example, partial amygdala or hippocampal kindling in cats increases defensive responding to rats (Adamec et al., 1976). There are also studies showing that long-term-kindling produces anxiogenic effect in rats (Adamec 1990, Kalynchuk et al., 2000). Further, kindled animals develop long-lasting deficits in declarative memory (Hannesson et al., 2003). With kindling progression, there is a gradual development of memory deficits assessed in a radial maze task (Sutula et al., 1995). Kindled rats also demonstrate retention deficits in the Morris water maze (Gilbert et al., 2000). Therefore, depending on the kindling protocol, electrical stimulations can disrupt a wide spectrum of memory functioning, from spatial memory and navigation to declarative and associative memory.

Cognitive deficits observed in animal models of TLE were confirmed in human patients. For instance, a recent longitudinal neuropsychological study documented that patients with TLE have pronounced intellectual decline after 15-30 years compared to age-matched controls tested with IQ (Jokeit et al., 2002; Oyegbile et al., 2004).

There is general agreement that cognitive dysfunctions produced by kindling are most likely caused by cumulative alterations in hippocampal structures with progressive reorganization. Hippocampal sclerosis, neuronal cell death, and mossy fiber sprouting are detectable morphologic consequences of the kindling application that eventually may lead to cognitive deficits. However, the exact mechanisms that control and mediate these cognitive deficits remain unknown.

Taking collectively, kindling can produce pathological changes on multiple levels, from cellular and morphological aberrations to behavioural outcomes. In the present experiment, we chose kindling as a model of epileptic seizures because of its advantageous opportunity to control the number and frequency of seizures over a given period of time. Importantly, kindling provides an opportunity to compare how seizures originating from different brain regions affect behaviour and cellular organization.

### **1.3. Neurogenesis**

During the past century, most of the neuroscientific community believed that adult neurogenesis was not possible. It was believed that once the mammalian brain reached maturity, the creation of new neuronal cells ceased. It has been nearly 50 years since the first report of neurogenesis in a mature rodent brain (Altman et al. 1963). It is now widely accepted that neurogenesis occurs in the human adult brain as well as in other vertebrates (Eriksson et al.,

1998; Spalding et al., 2013). Recent scientific progress provided compelling evidence that new cells are continuously added to the mammalian brain throughout adulthood (Dery et al., 2013, Spalding et al., 2013). The term “neurogenesis” was defined as a process of cell division and development from a stem cell into a mature fully functional neuronal cell (Ming et al., 2011). It should also be mentioned that developmental neurogenesis and the notion of adult neurogenesis significantly differ. This research line focused on the scope of adult neurogenesis.

Although it is true that most neurons are generated before birth and never regenerate, two brain regions in human brain play an avant-garde role creating new neuronal cells on a daily basis. They are the dentate gyrus (DG) of the hippocampus and subventricular zone (SVZ) of the lateral ventricles (LV). The rodent brain also demonstrates such activity in the olfactory bulb (Ernst et al. 2015). There are reports about additional sites involved in adult neurogenesis; however, we do not focus on them.

Adult neurogenesis involved in a broad spectrum of functions that are important for higher cognition, learning and memory (Cameron et al. 2015). Spatial coding, short-term memory and long-term memory formation, including its subdivisions of declarative and non-declarative memory, are supported by adult neurogenesis. Interestingly, studies also show an upregulation of neurogenesis following antidepressant treatment, indirectly, this may link depression and emotional regulation to adult neurogenesis (Hanson et al., 2011). We centre our research question on the connection between adult adult-born neurons and hippocampal-mediated memory functions (Aimone et al.2011; Deng et al.2010). There is a positive correlation between increased neurogenesis rates and enhanced memory performance on a Morris Water-Maze task; rat strains with low rates of neurogenesis appear to be slow learners (Kempermann et al., 2002). Conversely, ablation of neurogenesis may deteriorate performance on hippocampal-

dependent memory tasks (Clelland et al., 2009; Park et al., 2015). A recent study applying fast x-ray technology to arrest neurogenesis confirmed that adult-born hippocampal cells are required for novel object recognition, indicating that immature neurons get selectively recruited into hippocampal circuits during the update of stored information (Suárez-Pereira et al., 2015). Possibly, these processes provide a platform for pattern separation, which is critically important for learning processes. Pattern separation is a computational model explaining the ability to discriminate one memory from other stored memories. Importantly, ablation of neurogenesis affects cognitive abilities differently depending on the timeframe when the arrest of neurogenesis occurred. Selective inactivation of 4-week old cells, excluding those of other ages, significantly impairs retrieval of hippocampal memory (Gu et al., 2012). Possibly, this may indicate different functional roles of hippocampal cells during maturation. Clearly, the phenomenon of adult neurogenesis is a critical player in many domains of brain functioning.

Following the classic Pavlovian conditioning school, a trace fear conditioning paradigm may be considered as one of the manifestations of memory (Pavlov et al., 1927; Beylin et al., 2001). Mounting evidence demonstrates the active involvement of adult neurogenesis during all stages of memory process – formation, consolidation and storage of fear memories (Seo et al., 2015; Zhao et al., 2008; Snyder et al., 2005). There is a clear connection between trace conditioning tasks and neurogenesis. Several types of learning that depend on the hippocampal formation increase the number of newly generated neurons in the dentate gyrus. For example, learning a trace eyeblink conditioning task and spatial navigation in the Morris water maze increases the number of cells in dentate gyrus by enhancing their survival and proliferation (Gould et al., 1999a; Lemaire et al., 2000). Those tasks that do not require hippocampus do not influence on the process of cell growth (Gould et al., 1999). However, the connection between

neurogenesis and fear trace conditioning is not transparent. Published studies have shown that the suppression of neurogenesis can impair, enhance or even have no effect on trace conditioning tasks (Cuppini et al., 2006; Achanta et al., 2009; Jahol- Kowalski et al., 2009; Guo et al., 2011). Possibly, such obscurity may emerge due to methodological differences in neurogenesis suppression protocols. Taking into consideration the recent hypothesis that adult-born neurons may buffer hormonal and behavioural responses to stress, emotional factors may also modulate the results of such manipulations (Hayashi et al., 2008; Castilla-Ortega et al., 2011). Depending upon the type of conditioning protocol, the arrest of neurogenesis may deteriorate or improve performance on fear trace conditioning task (Seo et al. 2015). In those tasks that implement non-associative changes in fear expression, neurogenesis-arrested mice demonstrate elevation of fear behaviour. But in those procedures with minimal non-associative plasticity, neurogenesis-arrested mice show a decrease of fear expression (Seo et al., 2015). Therefore, it is important to emphasize that our study includes a trace fear conditioning protocol involving associative mechanisms.

Typically, any continuous process consists of multiple phases with its critical time-points and developmental stages, and neurogenesis is no exception to this principle. Adult neurogenesis follows a multifactorial pathway driving the stem cell into a mature neuron that eventually integrates into existing memory circuits. A cell generated from a basket cell flourishes into a mature neuronal cell, with its full functional capacity, within 4-6 weeks (Lledo et al., 2006). The process of cell growth goes through 3 consecutive stages: proliferation, maturation and survival. The adult rat brain creates thousands of cells every day, but most of them die within a few weeks (Brown et al., 2014). There are also data suggesting that adult neurogenesis may persist across the full lifespan from birth to physiological death (Knoth et al., 2010). Despite such promising

findings, it is clear that adult neurogenesis declines exponentially with age (McDonald et al. 2005). As mentioned above, the SVZ of the lateral ventricles and SGZ of the dentate gyrus as well as the olfactory bulb (OB) are restricted regions of the rat brain involved in processes of adult neurogenesis. In both places, dormant neuronal stem cells (NSC) give rise to neuronal progenitor cells (NPC) that have limitations in their proliferation. At this point, the fate of the new cell may be to become a glial cell or a neuron (Gage et al., 2000). The next phase of cell development is migration. Those cells designated to function in the olfactory bulb move from the SGZ via the migratory rostral stream to reach the olfactory bulb. The migration pathway of DG neurons is significantly shorter because it only involves having the cells move from the SGZ to the GCL. Finally, once a neuron migrates, it creates ramifications into the upper layers of the DG and connects to existing memory circuits.

Adult neurogenesis is regulated by numerous intrinsic and extrinsic factors. Among latter ones, well known are environmental enrichment, socialization and diet (Aimone et al., 2014). Voluntary running, as well as resistance training, increase the birth of new neuronal cells (Guerrieri et al., 2015), possibly by increasing vascularization and neurotrophic factors, providing more blood flow and oxygenation to hippocampal structures (Yak et al., 2014; Yarrow et al., 2010; Hötting et al., 2013). Interestingly, intermittent fasting in male rats also facilitates the proliferation of new neuronal cells (Manzanero et al., 2014; Lee et al., 2002). The above mentioned environmental factors coincide with memory enhancements tested on various behavioural paradigms (Kobilo et al., 2014).

Intrinsic factors also play an indispensable role in the regulation of adult neurogenesis. As mentioned, the local vasculature provides a platform for supplementing the process of cell growth. Internal layers of vascular tissue secrete factors that induce self-renewal of stem cells

and progenitor cell proliferation (Shen et al., 2004). Brain derived neurotrophic factor (BDNF) is another important contributing factor that has drawn much scientific attention recently. Studies indicate that high expression of BDNF is found within dentate gyrus in granule cells (Kato-Semba, Takeuchi, Semba, & Kato, 1997). Further, local infusions of BDNF into the hippocampus increases the proliferation of progenitor cells (Scharfman et al., 2005).

We also should denote the semantic difference between the words ‘regulation’ and ‘control’ because such a difference may create a misleading picture. The word ‘regulation’ in the literature is used to describe fluctuations from the baseline of neurogenesis. However, ‘controlling’ factors may produce a higher impact on the system because they are also involved in early embryogenic stages. The sequence of transcription factors such as Pax6-Neurog2-Tbr2-Neurod1-Tbr1 are a prominent example of a controlling factor (Kemperman et al., 2011). Collectively, all factors influencing adult neurogenesis may act in a synergetic fashion increasing or suppressing the process of cell maturation.

Using carbon method dating, there are data suggesting that the adult human brain can produce up to 700 new neurons every day. This represents 1.7% of neuronal cell turnover for every year of human life (Spalding et al., 2013). This is a significant event in human physiology and any alterations in the normal progression of neurogenesis may negatively contribute to many pathological conditions. Gaining a detailed understanding of the processes regulating adult neurogenesis may potentially bring new therapies to combat neurological and psychiatric diseases.

#### **1.4 Neurogenesis and pathology**

Adult neurogenesis has been associated with a number of psychiatric and neurological

diseases. Major depression, epilepsy, schizophrenia, anxiety and Huntington disease are well-described examples (Zhao et al., 2008; Schoenfeld et al., 2015; Gil-Mohapel et al., 2011).

Indeed, depression and chronic stress are associated with depletion of hippocampal neurogenesis (Jacobs, van, & Gage, 2000; Reif, Schmitt, Fritzen, & Lesch, 2007). It has been established that epileptic seizures increase the number of new neuronal cells (Parent et al., 2002). Different animal models of epilepsy confirm this phenomenon: pilocarpine induction, kainic acid and kindling all result in a higher number of proliferating cells. Our laboratory has shown that limbic kindling increases cell proliferation and promote dendritic hypertrophy (Botterill et al., 2015; Fournier et al., 2010). Indeed, there are numerous studies showing that seizures can distort adult neurogenesis through all stages of cell growth including proliferation, maturation, migration and integration (Parent et al., 2008). Thus, seizure-generated neurons display a significant difference in differentiation, migration, morphological maturation in contrast to neurons born under normal conditions. Seizures not only increase the number of new neuronal cells but accelerate the speed of their maturation (Overstreet-Wadiche et al., 2006). Additionally, to accelerated maturation, granule cells born after seizures also extend abnormal dendritic processes in the opposite direction – to the hilus (Shapiro et al., 2005; Ribak et al., 2006). These abnormal dendritic processes are known as hilar basal dendrites, and they are thought to contribute to hyperexcitability and may be considered as an epileptogenic hallmark (Pekcec et al., 2006).

Further structural reorganization includes mossy fiber sprouting, granule cell dispersion and hilar ectopic granule cell distortion, leading to a pathomorphological constellation that could facilitate hyper-excitation (Ribak et al., 2012). Despite such active scientific interest within the past few years, it is not clear yet what causes the acceleration in cell growth seen under conditions of repeated seizures. It is not clear what the functional relevance of increased neurogenesis might

be in the epileptic brain.

On a related topic, there appears to be a tight connection between adult hippocampal neurogenesis and status epilepticus, in that ablation of hippocampal neurogenesis before hippocampal-induced acute seizures in mice leads to a reduction in chronic seizure frequency (Cho et al., 2014) This study also shows a normalization of cognitive deficits associated with epilepsy. Further, these ameliorating effects are long-lasting and persist for up to 1 year. Another recent study suggests that after kainic-acid induced seizures in rats, swimming exercise may ameliorate cognitive deficits, even if the exercise is delayed (Geranial et al., 2016). Importantly, increased neurogenesis in TLE animal models is observed only in the acute phase. With the chronic progression of the disease, there may be a significant decline in hippocampal neurogenesis (Hattiangady et al., 2008; Danzer et al., 2012; Danzer et al., 2016).

Our laboratory previously showed that rats received 99 stimulations in the BLA region display a significant decrease in memory performance that coincides with significant increase of neurogenesis (Fournier et al., 2013). Thus, there is a link between aberrant neurogenesis and memory impairments in an animal model of epilepsy.

### **1.5. Markers of neurogenesis and cell activity**

For the past 20 years, significant technological advances opened the possibility of tracing the ‘private’ life of a neuronal cell and describing the factors influencing its birth, maturation and development. The new era of confocal imaging coincided with the discovery of histological markers such as BrdU, doublecortin and NeuN. These two events helped to further our understanding of factors influencing cell fate. During the cell growth process, different cell lineages create different cell types. They express exclusive cell surface proteins serving as a

marker to discriminate specific cell types and detect its stage of development. Here I selectively list and describe markers we applied in this study.

## **1.6 Neurogenesis markers**

Neuronal migration cell protein, also known as doublecortin (DCX), is a microtubule-associated protein expressed by neuronal cell precursors and immature neurons. Neuronal cell precursors express DCX continuously during the first 2-3 weeks of cell maturation. The decrease of its expression starts from the 2nd week (Brown et al., 2004). Adult-born neurons expressing DCX also display functional relevance despite their immaturity. A Recent study by Suarez-Pereira suggests that DCX-expressing neurons are necessary for updating memory in object-recognition tasks (Suarez-Pereira et al., 2015).

Bromodeoxyuridine (BrdU) is a synthetic nucleotide that mimics a thymidine structure. This unique molecular structure may substitute for thymidine during DNA replication and halt cell division in the S-phase. BrdU is a powerful instrument for monitoring the process of cell proliferation at any given time point. It is important to acknowledge that BrdU is a toxic and mutagenic substance. Thus, the dosage fluctuations may negatively affect the experimental procedure. BrdU is deemed to be a gold standard marker of cell proliferation and may help to visualize the process of cell mitosis at any given time point.

Nerve cells possess unique morphological structure allowing for the reception, transmission and storage of information. These operations may happen within a timeframe of milliseconds (neurotransmitter-mediated communication) or minutes like second messenger-mediated events. Trans-synaptic activation evokes a cascade of events including induction of new programs of gene expression (Black et al., 1987). Thus, numerous extracellular and

intracellular factors may modulate neuronal gene expression: membrane electrical activity, and neurotrophic growth factors like IGF-1 (Insulin Growth Factor -1 ) (Hu et al., 1994). Indeed, activation of immediate early gene (IEG) expression by extracellular stimuli is not a unique property of a neuronal cell. IEG's were first discovered in non-neuronal cells as a response to growth factors (Sheng et al., 1990). Expression of IEG's is transient and does not require protein synthesis, which may explain why their expression is so variable and fast. Nowadays, IEG expression is deemed to be a relevant molecular marker for a neuronal population that undergoes plastic changes that mediate learning processes (Kubik et al., 2007; Minatohara et al., 2016). It is also believed that IEG's are involved in transitioning the neuronal activity into long-term memories (Lanahan et al., 1998; Jones et al. 2001). As an example, pharmacologically induced convulsive and sensory stimuli provoke an upregulation of Arc, zif268 and Fos expression (Lyford et al., 1995). Behavioural tasks also induce significant IEG increases that are detectable in a variety of brain regions (Ramirez-Amaya et al., 2005).

### **1.6.1. FOS**

*c-Fos* is a proto-oncogene with the structure of an amino acid protein that acts inside the cell nucleus. Historically, the *c-Fos* family of genes were first discovered in rat fibroblasts as the transforming gene of the osteogenic sarcoma virus (Milde-Langosch 2005). The *c-fos* gene family was initially identified as an important factor in carcinogenesis (Mahner et al., 2008; Tulchinsky 2000). However, later it was established that Fos protein expression correlates with the level of a recent neuronal activity (Chung et al., 2005; Hoffman et al., 1991). For example, Fos protein expression in behaving rats reflects a summation of integration of activity-dependent calcium influx (Cruz et al., 2013).

*C-fos* acts rapidly in altering gene transcription in response to cell surface signaling (Dragoon et al., 1990) Expression of the *c-fos* gene peaks about 30-60 min after stimulation, and following a repression phase, it returns to its basal activity level (Greenberg et al., 1984). Fos is the protein product of *c-fos* and it may be induced by a variety of factors such as cytokines, tumor promoters, UV radiation and growth factors. FOS plays an important role in a variety of intracellular processes such as cell proliferation, differentiation and cell survival. Other studies suggest that Fos may also be involved in regulating the expression of BDNF and mediating mechanisms of neuronal excitability (Zhang et al., 2002).

### **1.6.2. ARC**

Activity regulated cytoskeleton-associated protein, or ARC (also known as Argo 3.1) is a protein that belongs to the immediate early gene family. Initially Arc was identified as the immediate early gene that was regulated by synaptic activity. This protein is involved in a variety of processes including embryonic development and the regulation of cell morphology, cytoskeletal organization (Liu et al., 2000). ARC may also play a prominent role in the acquisition and maintenance of several forms of hippocampal-dependent learning (Godowsky et al., 2001; Ploski et al., 2008). This immediate early gene is particularly implicated in the stabilization of LTP, LTD and homeostatic plasticity and may also contribute to modifying synaptic strength via postsynaptic density (Messaoudi et al., 2007). Importantly, Arc expression is critical for long term memory formation (LTM) and memory consolidation. Knockout Arc mice fail to demonstrate successful consolidation of fear memory, but acquisition and short-term memory formation were intact. (Plath et al., 2006). In a recent study, Arc knockout mice were presented as a unique animal model to study neural substrates of memory processing as the

animal demonstrates normal basal behavioural level while cognitive processing disrupted (cf. McCurry et al., 2010; Plath et al., 2006). Thus, the role of Arc is indispensable for healthy memory consolidation.

Interestingly, Arc gene expression is also related to functions that extend far beyond neuronal firing. Arc\Arg 3.1 regulates  $\beta$ -amyloid generation, which is actively involved in Alzheimer disease (Wegenast-Braun et al., 2009; Wu et al., 2011). In our study, we applied Arc antibody staining to detect recent neuronal activity following fear trace conditioning. This is a widely used effector protein and has maximal induction times of 5 minutes (RNA) or 60 minutes (protein) after behavioural testing (Botterill 2016). Further, there are data indicating that Fos and Arc signaling are necessary for the process of recruitment of newly generated cells for spatial learning and memory formation (Kee et al., 2007b, Trouche et al., 2007).

## **1.7. Neuroanatomy**

The next section explains the three brain regions that were involved in my experimental procedures. Each paragraph briefly describes the anatomical features, functional relevance and the connection to epileptogenesis as the prime focus of my research.

### **1.7.1. Hippocampus**

The hippocampus is a critically important brain structure for a proper memory functioning. It lies at a crossroads of all information processing in the brain receiving inputs from most of the cortex as well as from subcortical regions. Viewed this way, the hippocampus can be thought of as building connections to many brain structures directly or indirectly via the entorhinal cortex (Hebb 1949).

The hippocampal formation is a group of regions that includes the dentate gyrus, hippocampus, subiculum, presubiculum, parasubiculum and enthorinal cortex. The hippocampus anatomically has three pyramidal cell subfields: CA3, CA2, CA1 (CA means cornu ammonis); this organization appears to be similar across all mammalian species (Amaral et al., 2007).

The dentate gyrus is a trilaminar cortical structure that has a characteristic S shape. This is a cortical region representing an internal part of the hippocampal formation (Amaral et al., 2007). The granule cell layer (GCL) is a principal layer of the DG that comprises granule cells which are densely packed. A large portion of the GCL is located between CA1 and CA3. Three types of neurons represent a neuroanatomical organization of the DG: dentate principal granule cells, dentate pyramidal basket cells and mossy cells. The branch of granule cells extends through the molecular layer and ends at the ventricular surface. Granule cells give rise to unmyelinated axons called mossy fibers. The dentate gyrus is a principal anatomical input region for the hippocampal formation (Strien et al., 2009; Danzer et al., 2008; Lopez-Rojaz et al., 2016). It is widely accepted that the dentate gyrus has a primary function in providing a neurobiological platform for short-term memory and spatial orientation. It is deemed to be a pre-processing station for incoming information that most likely transforms the data by the principle of pattern separation. Thus, the dentate gyrus is capable of segregating similar information inputs and creating distinct differential patterns. This mechanism prepares incoming data for long-term storage in the CA3 region (Jonas et al., 2014).

Much of the scientific literature describes the hippocampus as a purely cognitive structure involved in the processes of memory functioning. Another portion of the literature treats the hippocampus as a regulator of emotion. Recent studies indicate different functional relevance within different subregions of the hippocampus (Fanselow et al., 2010; Schumacher et al., 2016)

This concept was initially proposed by anatomists, who observed differences in output\input projections between the dorsal hippocampus and ventral hippocampus (Swanson et al., 1977). Further studies also confirmed functional discrepancy, for instance, the dorsal hippocampus but not ventral hippocampus contributes to spatial memory processes. Thus, there was a rationale to segregate the hippocampus further into three zones: the dorsal, intermediate and ventral zones.

As I discussed in section 1.3, the dentate gyrus is a region that also carries an important function of adult neurogenesis (Piatti et al., 2013; Parent, 2007). Adult hippocampal neurogenesis considered to be a neuronal substrate for cognition, normal memory function and the process of memory ‘update’ (Suárez-Pereira et al., 2015; Anacker and Hen, 2017; Oomen et al., 2007). Apart from the memory processes, it has been already hypothesized that seizures may alter normal hippocampal neurogenesis (Parent 2002), which in turn may contribute towards epileptogenesis and cognitive impairments (Benardo, 2002; Cho et al., 2015).

A large body of epilepsy research has focused on the dentate gyrus because of its pivotal role in epileptogenesis (Dudek et al., 2007). The term epileptogenesis refers to the process by which previously normal brain is functionally altered and inclined towards the development of abnormal electrical activity that results in spontaneous recurrent seizures (Goldberg et al., 2013). There are several reasons why the dentate gyrus has generated a high level of interest for epilepsy research. The dentate gyrus is considered to be a critical structure in TLE because it is often presented as an epileptic focus and epileptogenic zone (Dudek et al., 2007). As previously discussed (Section 1.2), there is often neuronal cell loss following recurrent seizure attacks. Several concepts support the hypothesis that neuronal cell loss facilitates epileptic activity (Houser et al., 1996; Babb et al., 1991). Therefore, a reduction in the number of hilar neurons in

the epileptic brain may invoke network hyperexcitability because of the decrease in excitatory input to interneurons (Sloviter et al., 1987; Sloviter et al., 2003).

Another reason for scientific interest in the dentate gyrus stems from its unique trilaminar structure. Any synchronous activity in the laminar organization of the dentate gyrus generates large field potentials in the extracellular space (Dudek et al., 2007). A seizure may start in a nearby structure and it would be harder to detect abnormal electrophysiological activity because of the small field potential, but once synchronous activity involves the dentate gyrus, large amplitude field potentials or ‘fast ripples’ will appear on the electroencephalogram. These high-frequency electrographic oscillations serve as a hallmark of the epileptic activity (Bragin et al., 2002).

Collectively, the dentate gyrus is a highly important brain structure that is involved in processes such as memory, cognition, neurogenesis and epileptogenesis. Given my interests in the effect of kindling on cognition, I targeted the dorsal hippocampus as a kindling site to determine the impact of seizures on behaviour and neurogenesis.

### **1.7.2. Basolateral Amygdala**

The amygdala is an almond-shaped group of nuclei located deep and medially within the temporal lobes. This is a complex structure comprising about 10 different nuclei. The sensory input from various emotional states, predominantly fear and anxiety, enters the amygdala and transfers towards the basolateral complex (Kandel et al., 2000). The amygdala includes two major divisions that differ in their function and connectivity: the basolateral amygdala (BLA) and central amygdala (CeA). The basolateral complex (BLA) is further subdivided into the lateral (LA), basolateral (BL), and basomedial nuclei (BM). The central nucleus (CeA) is divided

into the lateral nucleus (CeL), the medial nucleus (CeM), and the intercalated cell masses (ICMs) regions (Duvarci et al., 2014). Glutamatergic neurons are the principal cellular component of the BLA and approximately 20% of the total cellular structure belongs to GABAergic interneurons (Pitkanen et al., 1995; Spampinato et al., 2011).

The amygdala has always been associated with the processing of fearful and aversive stimuli (Morris et al., 1996). In the early 1990's, a series of experiments with inactivation and surgical lesions of the BLA region suggested that the BLA plays a critical role in synaptic plasticity for the acquisition of Pavlovian fear learning and memory (Clugnet et al., 1990; LeDoux et al., 1990). Further, damage to the amygdala produces deficits in fear conditioning in humans (LaBar et al., 1995; Bechara et al., 1995). Conversely, fear conditioning is linked to increases in amygdala functional activity (Buchel et al., 1998; LaBar et al., 1998). Interestingly, there is also evidence suggesting that in addition to its role in fear and other negative reactions, the amygdala may also play a role in pleasurable reactions. For instance, lesions of the basolateral nuclei leave intact the learned association between the tone and the non-rewarding attributes of the food but disrupt the association between the tone and positive attributes of the food (Blundell et al., 2001; Kandel et al., 2000).

The amygdala has long been connected to the symptomatology of TLE. Stimulation of the amygdaloid complex may elicit the full spectrum of symptoms experienced by patients with intractable TLE (Gloor et al., 1982). The term 'dysphoric disorder of epilepsy' has also been proposed to describe a constellation of symptoms such as emotional instability, dysphoria, aggression and rage attacks accompanied by decreased volume of the amygdalar complex that suggests a causal link to the amygdaloid nucleus (Elst et al., 2009). There is also data indicating that selected populations of amygdaloid cell types demonstrate increased vulnerability to

epileptogenesis (Pitkanen et al., 1998). In fact, the amygdala of several species (including rats) is exquisitely susceptible to kindling (Goddard et al., 1969). Therefore, the basolateral amygdala is of great interest in relation to epileptogenesis and its influence on cellular and behavioural outcomes.

### **1.7.3. Caudate Nucleus**

The caudate nucleus (CN) is a distinct anatomical structure that acts in conjunction with the putamen (oval-shaped formation at the base of the forebrain) to form the dorsal striatum. The dorsal striatum in turn belongs to a larger structure called the basal ganglia. The CN receives projections from various structures; input projections are coming predominantly from all areas of the neocortex and one-tenth of afferent connectivity comes from the substantia nigra (SN) (Hokfelt et al., 1969). The functional role of the CN is usually described in the context of a functional relevance of basal ganglia pathways to which it belongs. Broadly speaking, it is implicated in the modulation of motor functions, sensorimotor coordination including response selection and initiation. However, the CN specifically has a function on its own, independently from the structures it is merged with. The CN plays an important role in stabilizing body posture and limb movement as well as regulating the speed and accuracy of directed movements (Villablanca et al., 2010). The frontal cortex acting with the CN are the functional part of the telencephalic inhibitory system; balancing posterior ventral-diencephalic and brain stem arousal sites (Villablanca et al., 2004).

The basal ganglia have never been associated with epileptogenic activity that leads to epilepsy. To date, there are no documented clinical reports that indicate a specific epileptiform

activity in this region and there are no specific changes following electrical activity in the after-discharge period.

However, it has been shown that the CN may cause a modulatory effect on hippocampal epilepsy (Vella et al., 1990). Possibly, the basal ganglia play an inhibitory role in TLE (Rektor et al., 2012). Some lines of research also demonstrated a significant reduction of seizure activity following caudate electrical stimulation. This effect was observed in different epilepsy models, cats suffering penicillin-induced hippocampal seizures (Rektor et al., 2012) as well as in cobalt-induced neocortical seizures (Grutta et al., 1988) demonstrated a significant reduction of seizure activity. Early clinical trials confirmed these initial observations and two patients had complete elimination of seizure activity following electrical stimulation of the CN; another four patients reported a significant reduction in seizure frequency (Sramka et al., 1976). Furthermore, intracaudate NMDA injections were capable of inhibiting pilocarpine-induced epileptic activity in rats (Turski et al., 1987). This study suggested that the caudate-putamen GABA-dependent efferent pathways may regulate the excitatory threshold of the limbic system. Thus, the CN can modify spiking activity of hippocampal structures and contribute to protective mechanisms against epileptiform activity. However, future research necessary to understand exact underlying mechanisms influencing electrical activity on the limbic structure.

### **1.8. Pavlovian Fear Trace Conditioning**

Classical Pavlovian fear conditioning involves learning an association between specific environmental stimuli and an aversive event. An animal learns to anticipate the emotionally significant event and mounts a physiological response to that event. The association of a conditioned stimulus with an unconditioned stimulus involves subcortical structures (Curzon et

al., 2009). However, if a context is the predictor of the aversive stimulus, then the hippocampus is involved in learning the association (Rudy et al., 2004). There are two types of fear conditioning protocols: delay and trace conditioning. In delay conditioning, the aversive conditioned stimulus (tone) co-terminates with the unconditioned stimulus (foot shock). In this scenario, the amygdala is primarily involved. Fear trace conditioning differs procedurally from 'delay' with the presence of an empty interval called a 'trace', which segregates the termination of a tone as a CS and the beginning of an aversive stimulus such as a footshock. There is additional complexity to trace conditioning due to the involvement of the hippocampus. Fear conditioning represents a behavioural paradigm that allows neuroscientists to study the neuronal circuitry underlying emotion, learning and memory. For decades, these paradigms have been extensively studied due to their simple experimental design and robust effects.

Several lines of research have demonstrated that the hippocampus and amygdala are critical for the acquisition of auditory fear trace conditioning. Lesions of the basal amygdalar nuclei produced deficits in auditory fear conditioning and contextual conditioning (Goossens et al., 2001). Further, rats with BLA lesions demonstrate a significant decrease in freezing behaviour in the presence of cues previously paired with footshock (Vazdarzhanova et al., 1998). The hippocampus, especially the dorsal hippocampus and CA3 region are critically important for the acquisition of trace fear conditioning (Curzon et al., 2009). Indeed, humans with damage to the hippocampus are able to acquire delay conditioning but not trace conditioning (Clark et al., 1998). Collectively, these studies provide support for the idea that auditory fear trace conditioning is both hippocampal-dependent and amygdala-dependent.

## **1.9. Experimental goal and the hypothesis**

The exact mechanisms that lead to cognitive deficits following persistent seizure activity remain unknown. One possible explanation is that aberrant adult hippocampal neurogenesis contributes to the memory impairments associated with epilepsy (Cho et al., 2015). Indeed, hippocampal neurogenesis plays important role in learning and memory (Eichenbaum, 1997), as discussed in the previous sections of Chapter 1. Newly generated granule cells demonstrate high structural plasticity in comparison to mature cells (Schmidt-Hiebe et al., 2004). It has been also demonstrated that seizures induce high rates of neurogenesis (Parent et al., 2002). Seizures distort not only the quantity, but also the quality of newly generated granule cells, because neurons born under seizure conditions display pathomorphological characteristics including exaggerated dendritic outgrowth and the maintenance of hilar-basal dendrites (Zhao et al., 2008). However, it is yet unclear whether seizure-induced changes in neurogenesis contribute to seizure-induced cognitive deficits. It is important to clarify this research question, because the research outcomes may potentially suggest a novel treatment approach for patients suffering from cognitive decline associated with TLE.

### **1.9.1. The specific aims and goals**

The specific aim of my thesis is to examine site-specific kindling outcomes on memory function, neurogenesis and rates of neuronal cell activation.

**Research question:** Do kindling induced changes in neurogenesis play a role in the cognitive deficits? To address this question, I will subject rats to long term kindling of three specific brain areas—the dorsal hippocampus, basolateral amygdala, and caudate nucleus. To examine

kindling-induced alterations in neurogenesis, I will inject BrdU at the time-point when electric stimulations impact hippocampal cell proliferation (30<sup>th</sup> electric stimulation). Kindling will continue after BrdU injections to the 99<sup>th</sup> stimulation, so that newborn cells can mature under conditions of repeated seizures. Then, the rats will be trained in a trace fear conditioning paradigm to examine the impact of site-specific kindling on fear memory retrieval. Shortly after behaviour testing, the rats will be sacrificed for immunohistochemical assessment of neurogenesis and neuronal cell activation via immediate early gene expression such as ARC and FOS.

**Hypothesis:** I hypothesize that kindling in different brain sites has differential effects on cognition, neurogenesis and levels of cellular activation. In particular, limbic but not non-limbic kindling would exaggerate neurogenesis and impair cognitive function of rats. If kindling induced changes in neurogenesis contribute towards cognitive deficits, then I expect that limbic, but not non-limbic kindling would alter neurogenesis and the expression of IEGs. Specifically, I expect that limbic kindled rats would have fewer newborn neurons expressing IEG's than non limbic or control rats, which would indicate impaired integration of newborn neurons after limbic kindling.

## CHAPTER 2

### **Seizures originating in different brain regions have differential effects on fear memory and the functional integration of seizure-generated granule neurons**

Nikita Nogovitsyn <sup>a</sup>, Justin J. Botterill <sup>b</sup>, Hector J. Caruncho <sup>c</sup>, Lisa E. Kalynchuk <sup>a</sup>

<sup>a</sup> Department of Medicine, University of Saskatchewan, Saskatoon, SK, S7N 5E5, Canada

<sup>b</sup> Department of Psychology, University of Saskatchewan, Saskatoon, SK, S7N 5A5, Canada

<sup>c</sup> College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, SK, S7N 5E5, Canada

## 2.1. Abstract

Epilepsy is known to instigate cognitive deficits. Acute seizures lead to an increase in the number of adult born neurons in the hippocampus (Botterill et al., 2015; Kokala et al., 2011). Recent findings demonstrate that there is a possible link between memory impairments and aberrant neurogenesis (Cho et al., 2015; Jessberger et al, 2015). Our previous studies established that the location where seizures originate may play critical role in the development of cognitive deficits (Botterill et al, 2017; Fournier et al., 2013). However, it is not known how specific seizure location impacts the rate of adult hippocampal neurogenesis. In present study, we subjected 25 rats to long-term kindling of various brain sites, then examined hippocampal neurogenesis and rates of neuronal activation following behavioural tasks. We labeled seizure-generated hippocampal neurons with BrdU after 30 stimulations and continued kindling until rats received a total of 99 stimulations. After the kindling procedure was complete, we tested rats on a trace fear conditioning task to examine the impact of seizures that originate from different brain sites on a fear memory retrieval. After fear conditioning, we sacrificed the rats and processed hippocampal tissue for immunohistochemistry to analyze neuronal activity in response to memory retrieval through immediate early gene expression. We found that long term kindling of limbic, but not non-limbic brain sites led to significant impairments in fear memory retrieval. Low rates of IEG activation in response to behaviour tasks were accompanied by exaggerated neurogenesis. These results suggest that seizures prevent the normal process of integration of newborn neuron into memory circuits. We also found that kindling of the dorsal hippocampus produced an extreme number of neuronal cells that corresponded to the lowest activation of neuronal cells and significant deficits in memory retrieval. Collectively, these data indicate that

seizure origin plays critical role in adult neurogenesis rates and provides more support for the idea that neurogenesis contributes to the cognitive deterioration that accompanies TLE.

## **2.2. Introduction**

Epilepsy is a group of neurological disorders encompassed by recurrent unprovoked seizures (Bernard et al., 2003). Temporal lobe epilepsy (TLE) is the most common form of epilepsy that is also complicated by pronounced cognitive deficits (Engel et al., 2001); it is notorious for its resistance to pharmaceutical intervention, thus to date there is no effective therapeutic strategy to target both seizure frequency and comorbid manifestations. Possibly, research endeavors should be directed towards the pathophysiology of epileptogenesis to answer clinical questions in regards to treatment of TLE.

For the past decade, much experimental attention has been directed towards understanding the phenomenon of adult hippocampal neurogenesis. It was established that new neurons are continuously added to hippocampal structures throughout the full life span of the mammalian brain. Adult neurogenesis is a multifactorial phenomenon regulated by a variety of internal and external factors such as diet, environment, stress and inflammation. There is also mounting evidence that acute seizures provoke an increase in the rate of cell proliferation, but the functional consequences of this increase is unclear. Further, there are numerous studies suggesting that neuronal cells produced after acute seizure attack may contribute to both epileptogenesis and cognitive impairments (Cho et al., 2015; Zhong et al., 2016).

Kindling refers to the gradual development and intensification of motor seizures resulting from daily delivery of electrical stimulations to specific brain regions. Kindling as a model has been successfully used to study epileptic seizures and its impact on neuronal plasticity. Our

previous studies have demonstrated that long-term kindling of the basolateral amygdala produces significant deterioration in fear memory retrieval that coincides with increased neurogenesis. Our findings supported the hypothesis that hippocampal neurogenesis is heavily involved in hippocampal-dependent memory tasks, thus perturbations in neuronal cell growth reflect on cognitive deficits. We also observed that seizures elicited through amygdala stimulation impair the normal activation of neurons during memory retrieval (Fournier et al., 2013).

In present experiment, we applied long-term kindling (99-stimulations) to three discrete brain regions to examine the impact of repeated seizures on hippocampal neurogenesis and neuronal activation in response to the retrieval of a fear memory. There were four treatment groups including the control group, caudate nucleus (CN) kindled, basolateral amygdala kindled (BLA), and dorsal hippocampus kindled (dHip). To examine the impact of repeated seizures on neuronal birth, we delivered three injections of BrdU following the 30<sup>th</sup>, 33<sup>rd</sup> and 36<sup>th</sup> kindled or sham stimulation. We continued the stimulations for another 4 weeks until rats received 99 stimulations and new neuronal cells marked with BrdU would reach maturity.

The next day, after the delivery of the last stimulation, rats were subjected to an auditory trace fear conditioning task. We quantified the expression of markers of recent neuronal activity such as Arc and Fos to examine levels of neuronal activity in response to behavioural testing following long-term kindling. To examine levels of activation of seizure-generated neurons that are 4 weeks old we quantified Arc positive cells on immunofluorescent labeling with BrdU marker.

A key aspect of this experiment is that all kindled rats experienced the same number of seizures, but those seizures were initiated in different brain regions. We expected that the BLA and dHip kindled rats would show cognitive deficits but the CN kindled rats would not. If

aberrant neurogenesis and deficient integration of newborn neurons is involved in these cognitive deficits, we should see these effects in the BLA and dHip kindled rat that show memory deficits, and not the CN kindled rats that do not show memory deficits. Our results largely supported this hypothesis, and confirm that seizure-generated neurons are unable to integrate into existing memory circuits.

## **2.3. Materials and methods**

### **2.3.1. Subjects**


We used 25 Male Long-Evans rats that were purchased from Charles River (Canada). Each rat weighted 200-250g upon arrival from the supplier. They were housed individually in a rectangular polypropylene cage with standard laboratory bedding and free access to food and water. The housing room was maintained at a temperature of 20<sup>0</sup>C ( $\pm$ 1 °C) on a 12:12 light:dark cycle with lights on at 8:00 am. All experimental procedures were performed during the light period of the light:dark cycle. All rats were treated in accordance with the Canadian Council for Animal Care and the University of Saskatchewan Committee on Animal Care and Supply. All efforts were made to minimize the number of animals used.

### **2.3.2. Surgery**

All rats were handled daily to acclimatize them to the animal colony room. After 1 week of acclimation, the rats were individually anesthetized with isoflurane (5%) and received an injection of a preoperative analgesic (Anafen, Ketoprofen, 10mg/kg, s.c.) to reduce pain and inflammation. After secure fixation in a stereotaxic apparatus, a mixture of isoflurane and oxygen was delivered through mouth tube to maintain anaesthesia. Isoflurane gas was

administered in a concentration of a 5% as initial step and 2.5% as a maintenance concentration. Targets of the surgery procedure were 3 discrete brain regions: the dorsal hippocampus (dHip), basolateral amygdala (BLA) or caudate nucleus (CN). A single bipolar electrode (MS-303-2-B-SPC, Plastics One, Roanoke, VA, USA) was implanted into left hemisphere of each rat in one of the relevant brain regions. The following stereotaxic coordinates were used: CN (+ 0.2 mm anteroposterior, + 3.2 mm mediolateral, - 5.7 mm dorsoventral); BLA (- 2.8 mm anteroposterior, + 5.0 mm mediolateral, - 8.5 mm dorsoventral); dHip (- 3.5 mm anteroposterior, + 2.6 mm mediolateral, - 3.1 mm dorsoventral) (Paxinos & Watson, 1998). To secure the electrode to the skull, 4 jeweler screws and dental acrylic were added as a final step of the surgery procedure.

All electrode placements were confirmed at the end of the experiment during the immunohistochemistry procedure. Sections containing electrode tracks were mounted on slides and used to confirm electrode placements. In addition, a Nissl stain was performed on selected brain sections to visualize the precise electrode location (Figure 1).



**Figure 1.** Representative photomicrograph of Nissl stained brain tissue. Electrode trace in dorsal hippocampus region - A; B – basolateral amygdala; C – caudate nucleus electrode mark. 1X magnification, Scale bar 100um.

### 2.3.3. Kindling

Following the surgical procedure, a 2-week recovery period took place in order to facilitate recovery. The rats were randomly distributed into 4 treatment groups: dHip (n=6), BLA (n=7), CN (n=6) and Control (n=6). The control group included rats who had electrode implants from each region of interest (dHip, BLA, CN).

Histological analysis revealed that 25 rats had correct electrode placements in the dHip, BLA and CN regions respectively. Representative electrode placements for each group are shown in Figure 1. Kindling progressed normally in all rats. Electrical kindling involved the daily application of a 1-second train of biphasic square wave pulses delivered at a frequency of 60Hz, an amplitude of 800  $\mu$ A (peak-to-peak), and a duration of 1 ms. Rats received 3 stimulations per day, 5 days per week with a minimum 3 hours break between each consequent stimulation. The total number of stimulations for each treatment group was 99-stim that was delivered approximately within 6.5 weeks. Rats assigned to the control group received 99 sham stimulations. This kindling protocol is identical to one used repeatedly in our laboratory (Fournier et al., 2013).

The behavioral convulsions induced by each electrical stimulation were scored using the revised 8 class extension (Pinel & Rovner, 1978) of Racine's original five class convulsion scale (Racine, 1972). Each numerical class of convulsions corresponds with severity of body motor involvement: Class 1: orofacial automatisms, Class 2: orofacial automatisms with head nodding, Class 3: unilateral forelimb clonus, Class 4: rearing with bilateral forelimb clonus, Class 5: rearing with bilateral forelimb clonus followed by falling, Class 6: multiple class 5 convulsions and falling episodes, Class 7: previous classes with running fit, Class 8: previous classes with intermittent muscle tonus. Based on this scoring scheme, a rat was considered to be "kindled"

when it displayed 3 consecutive class 5 convulsions. (Pinel & Rovner, 1978). There were no spontaneous seizures observed at any point during the experiment.

#### **2.3.4. BrdU injections**

One purpose of this experiment was to investigate development and maturation of seizure-generated neurons following the kindling procedure to determine whether pathologically born cells can integrate into memory circuits. Previous studies from our laboratory demonstrated that amygdala kindling increases the rate of cell maturation after 30 stimulations (Fournier et al., 2010). Therefore, in this experiment 3 BrdU injections at the dose of 75 mg/kg, 169 mg/m were delivered intraperitoneally 4 hours after the delivery of 30, 33 and 36 stimulations. All rats including the control group received 3 BrdU injections approximately 2.5 weeks after the first day of kindling procedure. Each injection was separated 24 hours apart. BrdU (BA-5002; Sigma-Aldrich) was dissolved in warm physiological saline solution (50 °C) and sterile filtered. After the last injection occurred, the kindling procedure resumed.

#### **2.3.5. Fear conditioning**

After the delivery of last stimulation, a number of behavioural tests took place in order to assess the cognitive consequences of kindling. Behavioural procedures comprised 4 consecutive days of training and testing:

Day 1. Habituation. Rats went through a habituation process to acclimatize them to the operant chambers. Two rats at the same time were brought to the brightly lit experimental room and then placed individually into operant chambers. They were allowed to explore the chamber for 10

min. Acetic acid solution 0.6% (v/v) was applied to eliminate olfactory cues for every rat going through operant chamber.

Day 2. Training. Rats were trained to learn the conditioned association. They were placed into the same room as they were on Day 1. After 180s of acclimatization, each rat received 7 sound tones (conditioned stimuli – CS) paired with a footshock as the unconditioned stimulus (US). Trace conditioning comprised a 16-s tone (85 dB, 2 kHz, 5 ms/rise fall time) followed by trace time of 30-s. A foot shock (0.9 mA) with a duration of 2-s was delivered at the end of the 30 sec trace period. Between every CS-US onset there was 180-s pause. The chambers were cleaned with acetic acid solution after every rat.

Day 3. Tone test. The tone test assessed the rat's ability to recall the association between the tone and the footshock. Rats were exposed to a novel operant chamber and received the same auditory tone from day 2. The tones were delivered 4 times (16 s duration, 85 dB, 2 kHz, 5 ms/rise fall time) with an interval of 198-s. Here we scored freezing time in response to the tone.

Day 4. Context test. The context test assessed the rat's ability to recall the association between the environment and the footshock. Rats were tested in the original environment that they were initially trained and habituated in. Rats were placed into the operant chamber for 480-s, and freezing was assessed during this time.

Freezing was defined as the absence of body movements except for those necessary for breathing. To evaluate defensive behaviour, we expressed freezing time as the percentage of time

spent freezing. The freezing episode of each rats was summed and divided by the number of observations then multiplied by 100.

### **2.3.6. Perfusions**

Two hours after the context test, the rats were deeply anesthetized with sodium pentobarbital (240 mg/kg, i.p.) and perfused transcranially with saline solution and then with cold (-20 °C) formaldehyde fixative (pH = 7.4). The brains were extracted and stored in the same solution until sectioning. Using a vibrating microtome (Vibratome 3000, Vibratome Company, St. Louis, MO, USA), all brains were sectioned at a thickness of 30 µm. Brain sections were securely stored in polycarbonate membrane dishes with a cryoprotectant solution, consisting of 30% (w/v) sucrose, 1% (w/v) polyvinylpyrrolidone, and 30% (v/v) ethylene glycol in 0.1 M PBS (pH = 7.4).

### **2.3.7. Post-mortem analysis: immunohistochemistry procedures**

We used standard immunohistochemistry protocols to assess the effect of kindling on the expression of several markers of neurogenesis and cell activation. All immunostaining procedures were performed with a free-floating technique to reach maximal antibody penetration.

#### **2.3.7.1. BrdU staining**

BrdU immunohistochemistry was used to visualize the number of proliferating cells. Sections were treated with 0.3% H<sub>2</sub>O<sub>2</sub> for 30 minutes. After TBS rinsing, the sections were gently transferred to 2 N HCl 267 at 45 °C for 1 hour to denature the DNA and expose the BrdU

antigen. 0.1M Borate Buffer was applied for 30 minutes to stop the chemical reaction and the tissue was then placed into blocking solution comprised of 5% NHS, 0.3% Triton-X-100 and TBS. After blocking, the sections were incubated with a primary mouse anti-BrdU monoclonal antibody (1:500, 48 h, 4 °C, Roche 272 Diagnostics) followed by incubation of the secondary antibody (biotinylated horse anti-mouse, 1:500, 2 h, room temperature, Vector Laboratories). Next, sections were placed in an avidin–biotin peroxidase complex for 1 hour (1:200, room temperature, Vectastain ABC Elite, Vector Laboratories). The last step of the immunolabeling procedure was visualization with 0.033% (w/v) 3,3'-diaminobenzidine (DAB) and 0.00786% 278 (v/v) hydrogen peroxide diluted in PBS until brown coloration of the tissue occurred. After the procedure, all sections were mounted onto slides and coverslipped the day after.

#### **2.3.7.2. Fos staining**

Fos protein immunohistochemistry was used to visualize neuronal activity after fear memory retrieval. Sections were exposed to hydrogen peroxide for 30 min in a concentration 0.3% at room temperature to stop endogenous peroxidase activity, followed by block-cocktail of 5 % (v/v) normal goat serum, 1 % (w/v) bovine serum albumin, and 0.3 % (v/v) Triton-X-100 in 0.1 M PBS for 1 hour. For the next 72 hours, all sections were incubated in primary Fos anti-rabbit polyclonal antibody at a concentration of 1:15,000 (Calbiochem, La Jolla, CA, USA) diluted in blocking solution. Next, sections were placed into biotinylated goat anti-rabbit IgG secondary antibody for 2 hours (1:500, Vector Laboratories, Burlingame, CA, USA) diluted in 0.3% (v/v) Triton-x-100 with 0.1M PBS, then treated with avidin-biotin-peroxidase (1:500, Vectastain ABC Elite, Vector Laboratories). Nickel-intensified DAB that comprised of 0.02% (w/v) 3,3'-diaminobenzidine, 2.5% nickel ammonium sulfate and 0.000083% (v/v) H<sub>2</sub>O<sub>2</sub> in

0.175 M sodium acetate visualized the immunolabeling procedure. After all rinses were done, sections were mounted onto glass slides and coverslipped.

### **2.3.7.3. Arc staining**

Arc protein immunohistochemistry was also used to visualize neuronal activity. Arc immunostaining started with exposing the sections to 0.3 % (v/v) H<sub>2</sub>O<sub>2</sub> for 30 minutes then transferring them into a block solution as described above. Primary ARC antibody (1:1000 rabbit anti-arc Synaptic Systems, Goettingen, Germany) diluted in block solution was applied to the sections for 48 hours. Next step was incubation in secondary antibody (1:500, 2 hours, Vector Laboratories) mixed with block solution. Immunoreaction was visualized using the DAB (0.05%) method (Botterill et al., 2015; Shu et al., 1988). Sections were then mounted onto glass slides and coverslipped.

### **2.3.7.4. Immunofluorescence double labelling**

Immunofluorescence double labeling was processed to visualize the presence of BrdU with Arc. This would identify newborn cells that were active during the recall of a fear memory. Following rinsing procedures, sections were treated with 2 N HCL for 30 min under 45 °C in a hot bath. Next, borate buffer was applied to stop the reaction and the section were washed in TBS 3 times. Prior to incubation with primary antibody, the sections were immersed in blocking solution (5% NGS, 0.3% Triton-X-100, 1% BSA in 0.1M TBS for 30 min). Following blocking procedure, the sections were incubated simultaneously for 48 hr at 4°C with Arc (1:500 anti-rabbit arc, Synaptic Systems, Goettingen, Germany) and BrdU (1:500 anti-mouse BrdU, Roche Diagnostics GmbH, Mannheim Germany) diluted in blocking solution. Further, the sections were

incubated in secondary antibodies: Alexa 488 Goat Anti-Mouse (1:500; Invitrogen, CA, USA) and Alexa 568 Goat Anti-Rabbit (1:500; Invitrogen, CA, USA) diluted in 5% NGS, 1% BSA, 0.1% Triton X-100 and TBS. This secondary cocktail was incubated for 3 hours at room temperature under minimal light exposure. After final rinsing, the sections were mounted onto charged glass slides and coverslipped with Citifluor fluorescent mounting medium. The final product of immunofluorescent staining was stored in minimal/no light exposure conditions at 4°C until further quantification analysis.

#### **2.3.7.5. Doublecortin immunohistochemistry**

Doublecortin (DCX) immunofluorescent labelling was used to confirm that BrdU labeling reflected proliferating neurons. Immunofluorescence with DCX was processed with the following protocol. Sections were rinsed with TBS solution (3 times for 10 min) and transferred to 5% H<sub>2</sub>O<sub>2</sub> diluted in 0.1 TBS for 30 min. Then, the sections were rinsed with TBS and incubated in sodium citrate at 85°C for 30min. Following TBS rinse, the sections were transferred to a blocking solution for 30 min (5% normal donkey serum, 1% Triton-X, 1% BSA dissolved in TBS). Next, the sections were incubated in primary DCX antibody for 48 hours at 4°C (1:250 Goat Anti-doublecortin c-18 sc-8066, Santa Cruz Biotechnology, CA, USA).

Following primary antibody exposure, the brain sections were rinsed with a TBS solution 3 times to prepare for secondary antibody exposure. Finally, sections were incubated in secondary antibody Alexa 488 Donkey anti-goat (1:250; Invitrogen, CA, USA) diluted in 5% NGS, 1% BSA, 0.1% Triton X-100 and TBS. The incubation procedure with secondary antibodies was 3 hr at a room temperature; samples were covered with tin-foil in a room with

ambient light. After a final rinsing procedure, the sections were mounted onto glass slides and coverslipped with citifluor fluorescent mounting medium.

#### **2.3.7.6. Quantification**

All quantification procedures were done by researchers blind to the treatment groups. Electrode placements were confirmed on immunostained tissue. Sections were examined using a Nikon Eclipse E800 microscope equipped with stage controller and digital camera (MicroFire, Optronics, Goleta, CA, USA). The quantification procedure involved three different methods: stereology counting, profile counting and manual counting.

**Stereology counting.** We counted BrdU positive cells that were visualized with DAB using unbiased stereological counting. BrdU positive cells were counted at a magnification of 40X in two distinct regions of the dentate gyrus: the subgranular zone (SGZ) and granule cell layer (GCL). The SGZ was identified as the region within a two-cell width in between the inner granule cell layer and the hilus. All subregions of interest were traced using the stereology software program StereoInvestigator (MicroBrightfield, Williston, VT, USA). Both ipsilateral and contralateral sides were included into the quantification. I counted cells in 5 brain sections per animal. All sections were 30  $\mu\text{m}$  thickness and sampled from every 12<sup>th</sup> brain section sectioned on the Vibratome. For the counting process, we applied a modified optical fractioning method to reduce oversampling (Kuhn et al., 1997). The total number of BrdU positive cells was estimated with the following formula:

$$N_{\text{total}} = \sum Q^- \times 1 / \text{ssf} \times A(\text{x, y step}) / a(\text{frame}) \times t / h.$$

$\sum Q-$  is the number of counted cells;  $ssf$  is the section sampling fraction (1 in 12);  $A(x, y$  step) is the area associated with each  $x,y$  movement ( $10,000\mu m^2$ );  $a(\text{frame})$  is the area of the counting frame ( $3,600 \mu m^2$ );  $t$  is the weighted average section thickness;  $h$  is the height of the dissector ( $12 \mu m$ ) (Botterill et al., 2015; Fournier et al., 2010; Lussier et al., 2013).

**Profile counting** was applied to quantify the number of Arc and Fos immunoreactive cells visualized with DAB. Profile counts were conducted bilaterally across 5 sections per brain. We counted Arc and Fos immunoreactive cells in three principal hippocampal subfields including the granule cell layer, subgranular zone, and hilar zone. First, at a magnification of 4X, we traced the boundaries of each hippocampal subfield using the rat brain atlas as a guide (Paxinos & Watson, 1998). Then, at a magnification of 40X, we counted cells using a meander profile counting method (Botterill et al., 2015; Knapska and Maren, 2009). The meander scan function is an integrated module of the StereoInvestigator software. The meander scan is a semi-automated scanning method that allow the user to count detected cells within a previously traced contour. And finally, to assess the density of immunoreactive cells, we divided the total number of immunoreactive cells by the total area measurement of the region of interest. The results for Arc and Fos cell counts are expressed as the average number of cells per  $mm^2$ .

**Manual counting.** All sections were captured using an epifluorescence microscope with a built-in stage controller and digital camera (MicroFire, Optronics, Goleta, CA, USA). Immunofluorescent pictures were captured under ambient light at a magnification of 4X under two distinct light spectrums to visualize BrdU (green light spectrum) and Arc (red light spectrum) positive cells separately. Both ipsilateral and contralateral sides of the brain were included in the quantification. BrdU and Arc positive cells were counted manually within the

SGZ and GCL subregions. Only cells that fluoresced brightly within the appropriate channel were included into the analysis. Manual counting was processed in three stages.

First, all BrdU positive cells within SGZ and GCL were counted and averaged for every animal. Second, all Arc positive cells were counted within the SGZ and GCL regions and then averaged. The total number of cells for every section was averaged for each animal. Importantly, the results of the manual counting for both BrdU and Arc corresponded with the number of cells counted using the profile and stereology counting methods. The third step examined the colocalization of BrdU and Arc. To evaluate the number of BrdU positive cells that also express Arc, I merged images from both fluorescent light spectrums using Photoshop. The brightness and contrast for each merged image was adjusted equally across treatment groups. Colocalized cells appeared as yellow-colored as a result of a convergence of red and green channels. All colocalized cells were counted within the SGZ and GCL regions in both hemispheres of the dentate gyrus. The total number of colocalized cells (yellow) were divided by the total number of BrdU positive cells (green) and then the acquired numbers were multiplied by 100 in order to express the data as a percentage.

#### **2.3.7.7. Statistical analyses**

All the data were analyzed using the Statistical Package for Social Sciences v21. Statistical significance was determined by one-way analysis of variance (ANOVA).

For each ANOVA, I considered a number of assumptions including the assumptions of homogeneity of variances and equality of means. All data were tested to confirm/reject the assumption of homogeneity of variances with the Levene test ( $p < 0.05$ ). The Welch's test was also applied to test the assumption of equality of means ( $p < 0.05$ ). If both assumptions were

confirmed, then the Tukey post hoc test was used to detect statistical differences among the groups. If the Levene or Welch's test indicated a violation of the assumptions ( $p < 0.05$ ), then I applied a non-parametric Games-Howell post hoc test (Toothaker, 1993) to detect statistical differences among the groups. The Games-Howell test is an appropriate alternative to the Tukey post hoc, because it does not assume equal variance and sample sizes (Ruxton & Beauchamp, 2008).

The statistical significance for all comparisons was set at  $P < 0.05$ . All graphs were built via Prism 6.0e where graph bars demonstrate the mean  $\pm$  standard error.

## **2.4. Results**


### **2.4.1. Kindling impacts fear trace conditioning**

Figure 2 shows the results of the behavioural testing. To evaluate baseline activity during day 1 habituation, we measured the number of midline crosses (see Figure 2A). There was no significant difference across groups in midline crosses ( $F(3,21)=1.466$ ,  $p=0.252$ ), thus kindling in both limbic and non-limbic groups did not impair motor activity.

On the training day (day 2) there were no significant differences between treatment groups, either in freezing time before the conditioned stimuli ( $F(3,21)=0.449$ ,  $p=0.617$ ) or during the trace period ( $F(3,21)=0.609$ ,  $p=0.617$ ) (Figure 2B). There were also no group differences in post-shock freezing time as a reaction to 7 CS-US pairings ( $F(3,21)=0.843$ ,  $p=0.486$ ). Prolonged freezing time across all treatment groups indicates successful learning of the conditioning association. The results of the training day procedures indicate that kindling did not affect the acquisition of fear learning.

Day 3 aimed to assess fear memory via the tone test. Our statistical analyses revealed group differences in the percent of time spent freezing (CS:  $F(3,21)=18.401, p=0.000$ ; PS:  $F(3,21)=18.401, p=0.000$ ). Post hoc analysis of this main effects further revealed that the BLA and dHip kindled rats froze less than the CN kindled rats or the control rats, as shown in Figure 2C.

Day 4 or the context test day aimed to assess memory for the association between the environment and the footshock. The BLA and dHip-kindled rats had a smaller percentage of freezing than the control or CN-kindled rats ( $F(3,21)=11.547, p=0.000$ ) (Figure 2D). In addition, we counted a number of fecal boli upon the completion the context test. Statistical analysis of these data revealed that the limbic kindled rats left significantly fewer fecal boli in the operant chamber room compared to rats in the other two groups ( $F(3,21)=20.887, p=0.000$ ) (Figure 2E).




**Figure 2.** Kindling affects fear trace conditioning. A) Kindling has no effect on habituation. B) Kindling had no effect on acquisition of the hippocampal dependent fear memories. C) Limbic kindled rats freeze significantly less than control and caudate kindled rats. D) First context test evaluated through the time spent freezing and showed a significant decrease of time spent immobile in limbic kindled rats E) Limbic kindled rats confirmed behavioural differences with control and caudate kindled rats, dHip kindled and BLA kindled found to have lower number of fecal boli in the operant chamber. (CN – caudate nucleus; BLA – basolateral amygdala; dHip – dorsal hippocampus;  $*p < 0.05$ ).


#### 2.4.2. Stereology counting for BrdU as a marker of neurogenesis

Figure 3 shows the results of the BrdU immunohistochemistry. A one-way ANOVA followed by post hoc Tukey test revealed a significant increase in the number of BrdU positive cells in the dHip kindled rats in comparison to the rest of the treatment groups on both the ipsilateral and contralateral sides (ipsilateral  $F(3,21)=17.141$ ,  $p=0.000$ ; contralateral  $F(3,21)=11.868$ ,  $p=0.000$ ; combined  $F(3,21)=14.717$ ,  $p=0.000$ ).

However, caution must be exercised due to the absence of homogeneity of variances assumption; that was confirmed by the Levene Statistic (ipsilateral  $F(3,21)=5.179$ ,  $p=0.008$ ; contralateral  $F(3,21)=7.472$ ,  $p=0.001$ ) and complemented by the Welch equality of means test (combined  $F(3,10.686)=10.432$ ,  $p=0.002$ ). Thus, there is a heterogeneity of variances among the groups. We should also take into consideration the fact that each treatment group comprises small sample sizes and importantly they are unequal ( $n=6$ ;  $n=7$ ;  $n=6$ ;  $n=6$ ;). Collectively, these factors decrease our statistical power making the classical Tukey post hoc test too conservative and incapable to detect potential significant differences between all treatment groups and creating the risk of a Type II error. Therefore, it is appropriate to re-analyze the results using the Games-Howel post-hoc test, which is a rather similar approach to Tukey but does not assume equal variances and equality in sample sizes. Using the Games-Howel test, we found that both the dHip kindled and BLA kindled rats had significantly more BrdU positive cells than the CN kindled and control rats (dHip-control  $p=0.029$ ; dHip-Caudate  $p=0.035$ ; dHip-BLA  $p=0.081$ ) (See Fig. 4). Hence, the present results are congruent with previous reports from our laboratory (Botterill et al., 2015; Fournier et al., 2013).




**Figure 3.** Comparative photomicrograph demonstrates a significant increase of BrdU positive cells in the dHip kindled and BLA kindled rats. Scale bar: 200  $\mu$ m.



**Figure 4.** Kindling affected BrdU expression in the SGZ and GCL in both ipsilateral and contralateral brain sites. There was a significant increase of BrdU positive cells in dHip kindled and BLA kindled groups in comparison to the control and caudate kindled groups ( $*p=0.01$ ).

### **2.4.3. Doublecortin fluorescence**

BrdU is an established method, widely employed by neuroscientists to track the process of cell proliferation. However, BrdU can label any proliferating cell not only neurons. Therefore, I performed immunofluorescent labelling with the immature neuronal marker DCX to confirm that the effects I was seeing with BrdU reflected changes in neurons. I examined the morphologic structure of DCX expressing cells within the SGZ and granule cell layers. Morphological characteristics and the topographic distribution of DCX expressing cells in each group confirmed that the BrdU immunostaining labelled proliferating neurons in the dentate gyrus (See Fig.5).




**Figure 5.** Illustrative example of DCX immunofluorescence across treatment groups. The pattern of neuronal expression within subgranular and granule cell layers corresponds with BrdU immunostaining. DCX expression confirms that BrdU positive cells are proliferating neurons.


#### **2.4.4. Profile counting for Arc immunoreactivity**

Figure 6 provides illustrative examples of Arc positive cells among all treatment groups. Tukey host-hoc analysis detected a significant decrease of Arc expression in limbic kindled groups within SGZ/GCL (ipsilateral  $F(3,21)=27.225$ ,  $p=0.000$ ; contralateral  $F(3,21)=20.854$ ,  $p=0.000$ ). Also there was a similar pattern of Arc expression within hilar zone of hippocampus (ipsilateral  $F(3,21)=9.772$ ,  $p=0.000$ ; contralateral  $F(3,21)=7.634$ ,  $p=0.001$ ). We also found that SGZ/GCL expressed more Arc positive cells in comparison to hilar zone (SGZ/GCL ipsilateral total mean =71.942, contralateral=74.803; hilar ipsilateral total mean = 4.05, contralateral total mean =4.720).

This quantification analysis indicates that limbic kindling significantly decreases neuronal activation after retrieval of a fear memory (See Fig. 7).



**Figure 6.** Comparative photomicrograph demonstrates the pattern of Arc expression in the dentate gyrus of four treatment groups. Scale bar: 200  $\mu$ m



**Figure 7.** Effect of kindling on Arc expression following context fear memory retrieval. Profile counting was performed in SGZ and GCL layers (left) and hilar zone (right). Both the dHip- and BLA-kindled rats showed a significant decrease in the number of Arc positive cells in comparison to the control and CN-kindled group [ $*p \leq 0.01$ ].


#### 2.4.5. Profile counting for Fos immunoreactivity

Figure 8 provides example of Fos expression across all treatment groups. Figure 9 shows the effect of kindling different brain sites on Fos immunoreactivity in the granule cell layer, subgranular zone and the hilus.


Figure 8 reveals a substantial number of Fos positive cells in the control and CN kindled groups within the SGZ and granule cell layer. One-way ANOVA test indicated a significant group difference for both hemispheres in the SGZ/GCL subregions (ipsilateral SGZ/GCL:  $F(3,21)=12.517, p=0.000$ ; contralateral SGZ/GCL  $F(3,21)=15.659, p=0.000$ ). Tukey post hoc was appropriate because the assumption of heterogeneity of variance was satisfied (Levene test for ipsilateral and contralateral  $F(3,21)=2.571, p=0.81$  and  $F(3,21)=2.805, p=0.065$ ). The Tukey post hoc test indicated that the dHip- and BLA-kindled groups had significantly fewer Fos positive cells within SGZ/GCL compared to the control and CN-kindled groups (all p values  $<0.006$ ). There were no statistical differences between the limbic kindled groups in either hemisphere ( $p>0.5$ ). Therefore, limbic kindling significantly decreased Fos expression within SGZ/GCL layers (See Fig.9).

One-way ANOVA for Fos expression within the hilus revealed a significant difference between the treatment groups (ipsilateral hilus:  $F(3,21)=13.384, p=0.000$ ; contralateral hilus:  $F(3,21)=5.409, p=0.006$ ). However, the Levene test indicated a heterogeneity of variance ( $p<0.05$ ). The Games-Howell post hoc test was therefore applied to compare means within the treatment groups. Only one pair had a significant difference within the treatment groups (CN-kindled to BLA-kindled:  $p=0.029$ ; all other pairs -  $p>0.064$ ).

Although the statistical difference was not detected in Fos expression within the hilus in the contralateral hemisphere, there remained an observable difference between CN-kindled compared the rest of the treatment groups. We will discuss this observation later (2.5.).



**Figure 8.** Comparative illustration of FOS expression across the four treatment groups. Photomicrographs at 4X magnification. Scale bar equals 200µm.




**Figure 9.** Kindling impacts FOS expression in limbic-kindled rats. There were significantly fewer Fos positive cells in the limbic kindled groups in comparison to the control and caudate groups within SGZ/GCL [ $*p \leq 0.06$ ].


#### 2.4.6. Double labelling for BrdU and Arc

Figure 10 shows pattern of BrdU and Arc expression using immunofluorescence. An example of the BrdU+Arc colocalization is provided in Figure 11. I counted BrdU positive cells within the SGZ and granule cell layers. Then, I counted Arc positive cells within the same regions of the dentate gyrus. Finally, I counted colocalized cells (BrdU positive cells that also expressed Arc). Statistical analyses of the BrdU and Arc counts were consistent with my findings reported above. Specifically, limbic kindling significantly elevated BrdU expression within the SGZ/GCL zones in both hemispheres (ipsilateral  $F(3,21)=21.795$ ,  $p=0.000$ ; contralateral  $F(3,21)=12.787$ ,  $p=0.000$ ). Limbic kindling also significantly decreased Arc expression within the SGZ/GCL areas in both hemispheres (ipsilateral  $F(3,21)=16.592$ ,  $p=0.000$ ; contralateral  $F(3,21)=20.157$ ,  $p=0.000$ ). The results from these manual counts demonstrate a similar pattern of BrdU and Arc distribution across all treatment groups (See Fig.12). Therefore, the present dataset corresponds with the results I obtained with profile counting and stereological counting.


Figure 13 shows the results from the colocalization analysis. Statistical analysis using one-way ANOVA followed by Tukey post hoc tests indicated a significant difference between treatment groups (ipsilateral  $F(3,21)=7.313$ ,  $p=0.002$ ; contralateral  $F(3,21)=8.374$ ,  $p=0.001$ ). The CN-kindled and control group had significantly more colocalized cells than the BLA-kindled and dHip-kindled rats. These results demonstrate that limbic-kindling prevented proliferating neurons from Arc activation in response to behavioural testing. The results of my experiment support previous reports from our laboratory (Fournier et al., 2013).



**Figure 10.** Pattern of fluorescent immunoreactivity of BrdU and Arc in each group. The green spectrum visualizes BrdU positive cells; Red – Arc positive cells. There is a significant induction of BrdU in the limbic-kindled groups. BrdU elevation was paralleled by a significant decrease of Arc expression in limbic-kindled rats. Pictures at 4X magnification. Scale bar equals 200 $\mu$ m.




**Figure 11.** Illustrative example of merged images. The arrow indicates a colocalized cell in the granule cell layer. Pictures at 4X magnification. Scale bar equals 100µm.



**Figure 12.** The effect of limbic kindling on BrdU and Arc fluorescent immunostaining following manual counting. Limbic kindling exaggerated neuronal proliferation that is paralleled with decreased Arc expression. \* $p=0.000$

### Colocalization of BrdU+Arc - SGZ/GCL



**Figure 13.** Colocalization analysis of merged images in immunofluorescent double labelling. Caudate and Control subjects demonstrates a significant difference compared to limbic kindled group. \* $p=0.05$

## 2.5. Discussion

Five principal findings arise from the present study. First, kindling does not impair the acquisition of fear learning. Second, kindling of limbic brain sites (basolateral amygdala, dorsal hippocampus) but not non-limbic brain regions such as the caudate nucleus, significantly impaired the retrieval of fear memories on subsequent behavioural tests. Third, markers of behaviorally relevant neuronal activity markers Fos and Arc were significantly higher in the control group and CN-kindled group compared to the BLA- and dHip-kindled rats. Fourth, limbic-kindled rats demonstrated significantly higher induction of hippocampal neurogenesis, indicated by BrdU and DCX. And finally, exaggerated hippocampal neurogenesis in limbic-kindled rats coincided with cognitive deficits.

In this study, we investigated the impact of long-term kindling in three discrete brain regions on neuronal activation of newborn granule cells. This neuronal activation was measured in response to the recall of a previously acquired fear memory. In line with a previous report from our laboratory (Fournier et al., 2012), we show that seizures increase the survival and accelerate the maturation of newborn granule cells. Limbic kindling has a negative impact on cell migration and its further integration in existing memory circuits.

Temporal lobe epilepsy affects several domains of cognitive functioning. Memory is comprised of several stages: acquisition, consolidation and retrieval. Memory can be further modified by the process of reconsolidation. In the present study, we demonstrated that seizures originating from limbic structures (BLA and dHip) significantly impaired retrieval of fear memories. Previous studies have shown that trace fear conditioning (auditory fear conditioning) is a form of episodic memory that unlike delayed conditioning is sensitive to lesions of the hippocampus. In our experiment, we show that long-term limbic kindling disrupts conditioned

fear in both the tone test and context test. However, the important observation that stems from the behavioural testing is that kindling did not impair the acquisition of hippocampal dependent memory. These results support previous reports from our laboratory (Botterill et al., 2016). Long-term kindling specifically isolated a portion of memory functioning but not all aspect of the memory. This observation may provide a hint towards further understanding of the pathophysiology of memory impairments under the condition of repetitive seizure attacks. Current neurophysiological studies have partially explained the molecular basis of memory acquisition. At the cellular level, trace fear conditioning requires the activation of NMDA receptors (LeDeoux, 2000; Wanisch et al., 2005) that help to detect coinciding events. With the help of AP5 (selective NMDA antagonist) infusions into the amygdala, it was demonstrated that rats fail to acquire fear conditioning (Miserendino et al., 1990). In line with this concept, Langton et al. also performed a series of MK-801 injections (NMDA receptor antagonist) that prevented rats from acquiring fear memories, importantly for both young and old rats (Langton et al., 2007). On the other hand, systematic NMDA injections have been shown to increase cognitive functions (Hlinak and Krejci, 2002; Koek et al., 1990). Both surgical and pharmacological experimental interventions suggest that the NMDA-receptor system is critically important for the formation of memories, but not for the maintenance and/or retrieval of such memories (Constantine-Paton 1994; Izquierdo 1991; Izquierdo et al., 1993; Liang et al., 1993; Quartermain et al., 1994; Rickard et al., 1994). Kindling is known to disrupt the NMDA system (Auzmendi et al., 2009; Shen et al., 2011; Auzmendi et al., 2008). There are eight components of the NMDA system (NR1, NR2A-D, NR3A-B, etc.). NR2A and NR2B are particularly interesting in this context because they are predominantly expressed in the hippocampus (Loftis et al., 2003). Importantly, both NR2A and NR2B subunits return to baseline at the 12-hour time point

after the last electric stimulation (Pratt et al., 1993). However, there are other reports indicating a significant reduction in the NR2A subunit in the amygdala and limbic forebrain in rats subjected to amygdala kindling (Prince et al., 1995). Possibly these controversies could be explained by the fact that these studies had different kindling duration.

Our experimental design involved regular electric stimulations over a prolonged period of time that allowed for adaptive and compensatory mechanisms to come into effect. Given the experimental design of our present study, we may hypothesize that rapid adaptive events, the stabilization of NMDA receptors in particular, within the time-gap between the delivery of the last stimulation and first exposure to the behavioural task may have allowed for a normal acquisition of Pavlovian conditioning. In fact, I conducted a preliminary immunohistochemical analysis of NR2B and PSD-95 labelling in the kindled rats from this experiment but a densitometry analysis revealed no significant differences between the treatment groups. The rats were sacrificed five days after the last stimulation. Given the fact that NMDA receptors are fast acting in nature, it is possible that the five-day gap allowed for the stabilization of the system. This potential stabilization effect may have prevented the detection of changes on the densitometry analysis.

The NMDA receptor is involved in a wide variety of neuronal functions such as synaptic plasticity, learning, memory and epileptogenesis (Li et al., 2009; Newcomer et al., 2000; Fraska et al., 2011; Blair et al., 2008). NR2B-containing NMDA receptors may play a crucial role in the cognitive deterioration observed in epileptic patients. The present discussion of possible neurobiological mechanisms that occur following repetitive seizures only partially covers the complex system that governs the process of memory acquisition, retention and retrieval. Therefore, the neurological basis of this pathology remains to be understood.

The mechanism of seizure-induced cognitive deficits remains unknown. However, there are several possible explanations for our observation that limbic kindling impaired fear memory retrieval in our experiment. In the adult animal, status epilepticus causes neuronal loss in the CA1, CA3 and the dentate gyrus. The pattern of cell loss depends on the animal model being used (Nadler, 1981; Ben-Ari, 2001). In addition to cell loss, the neuronal system is totally reorganized at multiple levels, with mossy fiber sprouting being one example of aberrant neuronal growth of granule cell axons in the epileptic brain (Represa et al., 1987; Sutula et al., 1988).

Our experiment demonstrates that kindling has different effect on neurogenesis depending on the brain site where the seizure originates. This finding may contribute towards the process of clinical evaluation of the severity of the condition in a clinical setting. We found that kindling originating from dorsal hippocampus has the most pronounced effect on the rate of neurogenesis, in that our stereological analysis revealed the highest number of BrdU positive cells in the dHip kindled group. These increases in neurogenesis coincide with poor performance on hippocampal dependent memory tasks.

Although dHip kindling exaggerates neurogenesis and decreases IEG activation, CN kindling induces somewhat opposite effects. Indeed, previous experimental evidence indicated that the caudate nucleus has a modulatory effect on hippocampal epilepsy (Vella et al., 1991). Electrical stimulation or NMDA microinjections has been shown to decrease seizure severity (Turski et al., 1988). In fact, clinical studies confirmed that low frequency brain stimulation of the caudate nucleus was able to suppress epileptic activity and disrupt both focal and generalized epileptic discharges (Šramka et al., 1990). Interestingly, the results of our study following Fos

analysis indicated a slight increase of neuronal activity in comparison to the control group, albeit statistical significance was not observed. It is also worth noting that the ipsilateral hemisphere showed slightly higher Fos activation within the hilus. It is possible that there are hemisphere-specific changes in response to kindling.

Stereological analysis of BrdU immunohistochemistry also showed similar trends. This observation may indirectly confirm that CN kindling may potentially inhibit the negative impact of repetitive seizures on neuronal circuits.

Our observations in CN-kindled rats correspond with reports from other laboratories. For example, central lateral nucleus high-frequency electric stimulations in rats results in cognitive improvement in performance and learning on a visual object recognition task. These improvements also coincide with increased expression of Fos and Zif268 in the caudate, putamen, and hippocampal regions (Shirvalkar et al., 2006). GABAergic system activation is one of the possible mechanisms that modulate behavioural outcomes following caudate stimulation. For instance, high frequency electric stimulation of the CN in vivo in rats demonstrated a significant increase in extracellular GABA expression in the caudate nucleus (Hiller et al., 2007). Therefore, evidence indicates that the CN is involved in inhibitory neural processes (Spehlmann et al., 1977; Rakic et al., 1962; Hiller et al., 2007). These reports may suggest that electric stimulation of the CN may produce potential therapeutic effects for patients suffering from TLE associated cognitive deficits.

In sum, the results of present experiment indicate that kindling of the dHip region and BLA enhance neuronal cell proliferation in the dentate gyrus. Limbic kindling impairs fear memory retrieval and seizure generated neurons demonstrate low neuronal activation following fear memory retrieval. Kindling in the dorsal hippocampus and basolateral amygdala has the

most pronounced negative effects on neuronal cell activation. Kindling in limbic regions prevented newborn neurons from normal integration into existing memory circuits. In conclusion, we provided evidence that aberrant neurogenesis as a result of limbic kindling plays a contributive role towards cognitive deficits.

## CHAPTER 3

### 3.1. General discussion

The general goal of my thesis was to advance the understanding of the mechanisms underlying the behavioural comorbidities associated with TLE. This thesis attempted to gain knowledge about the potential structural and functional changes in the hippocampus following repeated seizure activity.

I hypothesized that kindling in different brain regions would produce differential effects on behavioural outcomes. I further hypothesized that site-specific kindling would produce differential effects on neurogenesis and rates of neuronal cell activation. To that end, I examined the effect of long-term kindling in three distinct brain regions – the basolateral amygdala, dorsal hippocampus and caudate nucleus. As the product of my research, I received three principal sets of data.

First, I examined behavioural outcomes in response to long-term kindling with seizures originating from limbic or non-limbic brain sites. The day after the last kindling stimulation, rats were subjected to a trace fear conditioning paradigm to assess the impact of repeated seizures on the acquisition and retrieval of associative memory.

Second, I examined cell proliferation and neurogenesis following repeated seizures using the markers BrdU and DCX.

Third, I examined the rate of neuronal activation in response to exposure to a pre-conditioned (aversive) environment following behavioural training.

I found that limbic kindling significantly impaired fear memory retrieval. Behavioural outcomes paralleled increased neurogenesis in limbic kindled rats. In contrast, neuronal activation rate was diminished after limbic kindling, and very few BrdU positive cells also

expressed Fos and Arc. These results suggest that limbic, but not non limbic kindling prevent seizure-generated neurons from properly integrating into existing memory circuits in the hippocampus.

### **3.1.1. Aberrant neurogenesis and cognition**

The addition of a large number of neuronal cells generated in response to kindling indeed impaired cognitive performance in rats. Colocalization analysis within my thesis work suggested that limbic kindling prevented newborn neurons from normal integration into the existing memory circuits. This hypothesis is supported by a previous report from our laboratory that seizure generated neurons are less likely to colocalize with the marker of recent neuronal activity (Fournier et al., 2013).

However, other researchers found evidence to support an alternative hypothesis. Granule-like cells produced after pilocarpine induced seizures appear to participate in the underlying neuronal activity that appears during chronic behavioural seizures (Sharfman et al., 2002). Possibly, neurogenesis in the dentate gyrus following pilocarpine seizures can lead to new cells that mature into functional neurons and integrate into the functional circuits (Sharfman et al., 2000, 2002). Importantly, the Sharfman laboratory did not examine the level of neuronal activation of those ‘recruited seizure newborn’ neurons in response to cognitive tasks. Therefore, although seizure-generated granule cells may mature into functional neurons, it is not clear that these post-seizure neurons can function properly in memory tasks.

There are other interesting perspectives that attempt to explain a connection between aberrant neurogenesis and cognitive deficits. Possibly, aberrant integration and the presence of many newly generated neurons impair memory performance by degrading network performance

(Fournier, 2009; Becker et al., 2005; Wiscott et al., 2006). The constant addition of newly generated neurons into a neuronal network is compensated by randomly chosen pre-existing neurons (Chambers et al., 2004). If indeed, seizure-generated neurons, with their pathomorphological characteristics, still succeed to integrate into a memory circuit, then another possible explanation is a ‘wash out’ effect. In other words, the permanent accumulation of granule cells simply creates a competition for new neurons to integrate into the network, in addition to this a constant turnover of new neurons create a constantly updating network that may impede normal memory retention (Frankland et al., 2013; Akers et al., 2014). To support this idea, there is evidence that elevated levels of neurogenesis promote forgetting of established memories (Akers et al., 2014).

### **3.1.2. Why do seizures impede neuronal activation?**

The results of my thesis work support concept that seizures dramatically decrease the neuronal activation that is necessary for normal memory functioning. Levels of neuronal activation were severely dampened in the limbic kindled rats. Although aberrant neurogenesis may play an important role in the cognitive deficits that result from kindling, this does not explain the mechanism of neuronal deactivation. It is unknown what factors might mediate this pathological process.

As it has been previously discussed (section 1.2), kindling induces pathological changes on multiple levels. These pathologic changes involve a myriad of events that are connected into the one unified synergetic system. This synergy makes it increasingly difficult to select and understand the leading factors that govern neuronal downregulation in response to recurrent

seizures. However, there are several possible mechanisms that may play dominant role in ‘neuronal deactivation’ within a memory network.

Recurrent seizures evoke a cascade of compensatory mechanisms. One of them is the observable strengthening of the GABAergic system. For instance, new cells that are born under the condition of status epilepticus receive increased inhibitory activity (Jakubs et al., 2006). Seizures accelerate the process of cell maturation (Overstreet-Wadiche et al., 2006) and at the same time seizure-generated neurons receive inhibitory inputs sooner than the newborn cells created under normal conditions. This pathologic acceleration possibly affects a critical period of neuronal cell development, when cells may exhibit increased plasticity. Therefore, seizure-generated neurons are ‘silenced’ by the GABAergic system before they unfold their potential of synaptic plasticity. Further, it is well documented that seizures cause mossy fiber sprouting (Blaabjerg et al, 2007; Danzer et al., 2009). It has been hypothesized that the terminals of those sprouting neurons may release GABA into the dendrites of granule cells (Buckmaster, 2012; Santhakumar et al., 2005). Therefore, mossy fiber sprouting may also help to ‘silence’ both newborn and mature neurons. Viewed this way, the reduced neuronal activation seen in the limbic-kindled rats in this experiment may be the manifestation of a compensatory mechanism. Thus, increased inhibitory input that helps to ‘resist’ further epilepsy progression comes at the general cost of neuronal deactivation.

Another possible explanation for the reduced neuronal activation following epileptic seizures comes from the perspective of long term potentiation (LTP). Long term synaptic plasticity is a key mechanism that allows for normal memory function and cognition (Malenka, 1994). LTP is associated with increased synaptic connectivity in response to high-frequency stimulation. The induction of LTP requires both GABAergic and glutamatergic systems (Davies

and Collingridge, 1993). Physiological high-frequency stimulation opens NMDA receptor channels allowing the influx of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$ . Inside the post-synaptic neurons,  $\text{Ca}^{2+}$  activates calmodulin kinase II and protein kinase C. These enzymes induce LTP that serves for the benefit of memory processes (Hamed, 2007). Electrophysiological studies in the rat hippocampus demonstrated that persistent seizure activity has a destructive effect on synaptic plasticity. Seizures cause widespread induction of LTP that reduce overall hippocampal plasticity (Reid et al., 1997; Hamed, 2007). This reduction of hippocampal plasticity following seizure activity may partially explain the limited neuronal activation in the dentate gyrus.

Over the past decade, there was a significant progress in understanding the pathophysiology of epilepsy. However, further scientific efforts are necessary to create a holistic understanding of epileptogenesis and the causes of the cognitive deficits associated with epilepsy.

### **3.2. Limitations**

Kindling is a well-studied animal model of epilepsy. Kindling has been known since the 1960's and has been extensively applied in various experimental situations. Experiments with electric stimulation have been reported in various vertebrate species. Importantly, the behavioural outcomes resulting from amygdaloid kindling are very similar across many species. Local electrical stimulations of the amygdala or the hippocampus are associated with a progressive increase in seizure susceptibility and severity. As it was previously discussed, kindling can induce neuronal degeneration in the limbic region that is very similar to those changes observed in patients with TLE. This observation strengthens the validity of kindling as a model of TLE. Further, kindling has a high predictive validity because anticonvulsant drugs that

successfully demonstrate therapeutic action against kindling are also effective in human patients. Interestingly, kindling also shares many similarities with the behavioural sensitization resulting from psychostimulant drugs (Pierce and Kalivas, 1997; Post and Weiss 1989).

Among other strong advantages of kindling there is a location specificity, thus researchers are able to elicit seizures in a specific brain site and demonstrate high control of the number of seizures and their frequency. This allows for the study of time-sensitive events that parallel epileptic seizures.

However, kindling has received a substantial amount of criticism within the past decade. Spontaneous epileptic seizures are the main feature of the epilepsy in human patients. However, kindling does not elicit spontaneous seizure activity. Although it is possible to elicit spontaneous seizures after more than 280 stimulations, kindling does not successfully mimic this clinical feature in its normal application. Another disadvantage of kindling is that it is still not clear whether kindling can induce epilepsy in a human brain. There are few consistent reports when patients received series of ECT stimulations and later progressed into full-blown epilepsy. In fact, there is a clinical case of a patient that experienced more than 1200 ECT sessions and never developed spontaneous recurrent seizures (Lipman et al., 1985).

Kindling as a model of epilepsy has remained mainstay of epilepsy research for more than 40 years. But the question remains about how the results from kindling experiments advance our understanding of the underlying neurobiology of human epilepsy. Therefore, kindling has a limitation in regards to clinical relevance.

Another limitation of my thesis work stems from sex specificity. Sex differences may play a role in kindling progression and its behavioural outcomes. Sex hormones and their neurosteroid derivatives exert regulatory activity on the excitatory properties of the hippocampus

(Smith et al. 2002). Our experimental procedures were conducted exclusively with male rats. I would like to acknowledge that this is one of the limitations of my present work. Estrogens, androgens and progestins are well-known to alter hippocampal electrophysiology.

Steroid hormones play key role in the neuroendocrine control of neuronal excitability and seizure susceptibility (Herzog et al., 2003; Verotti et al., 2007). Steroid hormones are synthesized in ovarian, gonadal and adrenal structures. In men, the dominant steroid hormones are testosterone and dihydrotestosterone; both exhibit anabolic and androgenic activity. In women, the main circulating reproductive steroid hormones are the estrogens and progesterone. Testosterone has been shown to decrease excitation threshold and increase seizure susceptibility. However, testosterone may exhibit a dual role on neuronal excitability, because it may produce both proconvulsant and anticonvulsant effects depending on the animal model and seizure type (Reddy et al., 2008). The proconvulsant activity of testosterone may be explained by its aromatization to estradiol. Both animal and clinical studies indicate that testosterone metabolites increase seizure activity (El-Khayat et al., 2003; Herzog et al., 2003). Conversely, the androgenic wing of testosterone has a protective role against seizures that were induced by kainic acid (Reddy et al., 2004; Frye et al., 1999). Therefore, testosterone may exert variable effects on seizure activity: proconvulsant at higher doses and anticonvulsant at lower doses. In clinical settings, testosterone injections have not been shown to improve seizures (Herzog et al., 1998). The female estrus cycle also causes fluctuations in hippocampal excitability, which may produce differential risks for seizure susceptibility at different points of the hormonal reporting period (Warren et al., 1995; Scharfman et al., 2006). Unlike testosterone, the estrogens generally facilitate seizure activity (Wooley et al., 2000). Seizures and hormonal activity have a bidirectional connection. Limbic kindling distorts ovarian hormone cyclicality in rats (Edwards et

al., 2000). The effect of the estrogens has been studied using animal model of epilepsy. Epileptic female rats exhibit elevated seizure frequency that coincides with peaking time points of estrogen expression during the ovarian cycle (Sharfman et al., 2008). The estrogens also demonstrate a systemic influence on neuronal functioning. Neurogenesis is no exception, as estradiol may rapidly upregulate neuronal proliferation in the dentate gyrus of female rats (Barha et al., 2009).

Given the limitations of my present work, it is important to note that future studies may further investigate the connection between neurogenesis and site-specific epilepsy under an experimental design that would consider sex differences and kindling limitations. Finally, most of the findings described in the present work are based on animal models, therefore caution must be exercised when extrapolating these findings to the human brain.

### **3.3. Future directions**

The results of my thesis further establish the importance of neurogenesis as a contributing factor in the cognitive deficits associated with TLE. These results may potentially impact a therapeutic strategy of future epilepsy treatment. Due to the importance of normal neurogenesis, it is logical to hypothesize that patients with TLE may benefit from therapeutic interventions that modulate adult hippocampal neurogenesis. For example, rapamycin has been already suggested as one possible option for pharmacologic intervention (Wong 2011; Canpolat et al., 2014). Rapamycin administration can reverse the cognitive deficits produced by status epilepticus in rats (Brewster et al., 2013). These effects of rapamycin may be mediated by a reduction of aberrant neurogenesis and decrease of mossy fiber sprouting (Butler et al., 2015; Zheng et al.,

2009). Therefore, future research should investigate the development of pharmacological modulators of hippocampal neurogenesis.

As mentioned in previous paragraphs, the development, maturation and migration of newborn neurons in the mature brain is mediated by a cascade of intrinsic factors, one of them being reelin (Teixeira et al., 2012). Reelin is a large extracellular matrix protein expressed predominantly by GABAergic interneurons in the hippocampus, and it has been shown to play a critical role in neuronal migration (Teixeira et al., 2012). Reelin-deficient mice have significantly fewer newborn granule neurons as compared to their control counterparts (Won et al., 2006). Conversely, intrahippocampal infusions of reelin enhance the cognitive abilities of mice; this enhancement parallels morphological reorganization of hippocampal structures (Rogers et al., 2011). Intrahippocampal infusions of recombinant reelin have also been shown to reduce neuronal aberrations following epileptic seizures in mice subjected to the kainite model of TLE (Müller et al., 2009). Recently, it was demonstrated that persistent seizure activity leads to a significant decrease in reelin expression, both in the human and animal brain (Frotscher et al., 2016).

Previous reports suggest that that reelin plays an essential role in adult neurogenesis. Considering the relationship between seizure-induced neurogenesis and cognitive decline in TLE, it is logical to hypothesize that reelin in the future reelin may become a therapeutic option for patients suffering from drug-resistant forms of epilepsy and cognitive deficits. Reelin has already been considered as a potential therapy for depression and mood disorders (Caruncho et al., 2016). Therefore, it is interesting to apply reelin as a therapeutic intervention in an animal model of epilepsy such as kindling. However, there has been criticism regarding safe and effective routes of reelin administration, as reelin is a large protein that has difficulties

crossing the blood brain barrier (Caruncho et al., 2016). Further experimental procedures delineating the pharmacodynamics and pharmacokinetics of reelin are needed. Currently, the field has a knowledge gap in understanding how reelin is affected by pathology, and how the supplementation of reelin could be efficacious in treating pathology. Further studies should apply these questions to animal models of TLE.

## REFERENCES

- Adamec R. Behavioural and epileptic determinants of predatory attack behaviour in the cat. 1975;2:457-466.
- Aimone JB, Li Y, Lee SW, Clemenson GD, Deng W, Gage FH. Regulation and Function of Adult Neurogenesis: From Genes to Cognition. *Physiol {Rev}*; 2014. p. 991-1026.
- Akers KG, Martinez-Canabal A, Restivo L, Yiu AP, Cristofaro AD, Hsiang H, et al. Hippocampal Neurogenesis Regulates Forgetting During Adulthood and Infancy. *Science* 2014 may;344(6184):598-602.
- Altman J, Das GD. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol* 1965 jun;124(3):319-335.
- Alvarez-Buylla A, Lim DA. For the long run: maintaining germinal niches in the adult brain. *Neuron* 2004 mar;41(5):683-686.
- Amaral DG, Scharfman HE, Lavenex P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Prog Brain Res* 2007;163:3-22.
- Anacker C, Hen R. Adult hippocampal neurogenesis and cognitive flexibility “ linking memory and mood. *Nat Rev Neurosci* 2017 jun;18(6):335-346.
- Auzmendi J, González N, Girardi E. The NMDAR Subunit NR2B Expression is Modified in Hippocampus after Repetitive Seizures. *Neurochem Res* 2009 may;34(5):819-826.
- Avanzini G, Depaulis A, Tassinari A, de Curtis M. Do seizures and epileptic activity worsen epilepsy and deteriorate cognitive function? *Epilepsia* 2013 nov;54 Suppl 8:14-21.
- Babb TL, Kupfer WR, Pretorius JK, Crandall PH, Levesque MF. Synaptic reorganization by mossy fibers in human epileptic fascia dentata. *Neuroscience* 1991;42(2):351-363.
- Shahram Bahramia, Finn Drabløsa. Gene regulation in the immediate-early response process. *Advances in Biological Regulation* 2016 sep;62:37-49.
- Bechara A, Tranel D, Damasio H, Adolphs R, Rockland C, Damasio AR. Double dissociation of conditioning and declarative knowledge relative to the amygdala and hippocampus in humans. *Science* 1995 aug;269(5227):1115-1118.
- Bechara A, Damasio H, Tranel D, Anderson SW. Dissociation Of Working Memory from Decision Making within the Human Prefrontal Cortex. *J Neurosci* 1998 jan;18(1):428-437.
- Becker S. A computational principle for hippocampal learning and neurogenesis. *Hippocampus* 2005;15(6):722-738.

Beckman D, Santos LE. The Importance of Serotonin in Exercise-Induced Adult Neurogenesis: New Evidence from Tph2<sup>-/-</sup> Mice. 2013 sep.

Belzung K. Neurogenesis and Neural Plasticity: Springer; 2013.

Benardo LS. Epileptogenesis and Neurogenesis Share the Same Genes. *Epilepsy Curr* 2003 sep;3(5):162-163.

Ben-Ari Y. Cell death and synaptic reorganizations produced by seizures. *Epilepsia* 2001;42 Suppl 3:5-7.

Johan Bengzon, Zaal Kokaia, Eskil Elmér, Avtandil Nanobashvili, Merab Kokaia, and Olle Lindvall .Apoptosis and proliferation of dentate gyrus neurons after single and intermittent limbic seizures. *PNAS* 1997 sep;94(19):10432-10437.

Bhalla D, Godet B, Druet-Cabanac M, Preux P. Etiologies of epilepsy: a comprehensive review. *Expert Review of Neurotherapeutics* 2011 jun;11(6):861-876.

Birbeck GL. Epilepsy Care in Developing Countries: Part I of II. *Epilepsy Curr* 2010 jul;10(4):75-79.

Black IB, Adler JE, Dreyfus CF, Friedman WF, LaGamma EF, Roach AH. Biochemistry of information storage in the nervous system. *Science* 1987 jun;236(4806):1263-1268.

Blair RE, Sombati S, Churn SB, DeLorenzo RJ. Epileptogenesis causes an N-methyl-d-aspartate receptor/Ca<sup>2+</sup>-dependent decrease in Ca<sup>2+</sup>/calmodulin-dependent protein kinase II activity in a hippocampal neuronal culture model of spontaneous recurrent epileptiform discharges. *Eur J Pharmacol* 2008 jun;588(1):64.

Blundell P, Hall G, Killcross S. Lesions of the basolateral amygdala disrupt selective aspects of reinforcer representation in rats. *J Neurosci* 2001 nov;21(22):9018-9026.

Botterill JJ, Brymer KJ, Caruncho HJ, Kalynchuk LE. Aberrant hippocampal neurogenesis after limbic kindling: Relationship to BDNF and hippocampal-dependent memory. *Epilepsy Behav* 2015 jun;47:83-92.

Botterill JJ, Fournier NM, Guskjolen AJ, Lussier AL, Marks WN, Kalynchuk LE. Amygdala kindling disrupts trace and delay fear conditioning with parallel changes in Fos protein expression throughout the limbic brain. *Neuroscience* 2014 apr;265:158-171.

Botterill JJ, Nogovitsyn N, Caruncho HJ, Kalynchuk LE. Selective plasticity of hippocampal GABAergic interneuron populations following kindling of different brain regions. *J Comp Neurol* 2017 feb;525(2):389-406.

Boylan LS, Flint LA, Labovitz DL, Jackson SC, Starner K, Devinsky O. Depression but not seizure frequency predicts quality of life in treatment-resistant epilepsy. *Neurology* 2004 jan;62(2):258-261.

Bragin A, Mody I, Wilson CL, Engel J. Local generation of fast ripples in epileptic brain. *Journal of Neuroscience* 2002;22(5):2012-2021.

Bramham CR, Alme MN, Bittins M, Kuipers SD, Nair RR, Pai B, et al. The Arc of synaptic memory. *Exp Brain Res* 2010 jan;200(2):125-140.

Brandt C, Ebert U, Löscher W.. Epilepsy induced by extended amygdala-kindling in rats: lack of clear association between development of spontaneous seizures and neuronal damage. *Epilepsy Res* 2004 dec;62(23):135-156.

Braun SMG, Jessberger S. Adult neurogenesis: mechanisms and functional significance. 2014 may.

Brewster AL, Lugo JN, Patil VV, Lee WL, Qian Y, Vanegas F, et al. Rapamycin Reverses Status Epilepticus-Induced Memory Deficits and Dendritic Damage. *PLoS One* 2013 mar;8(3).

Brown JP, Couillard-Despres S, Cooper-Kuhn CM, Winkler J, Aigner L, Kuhn HG. Transient expression of doublecortin during adult neurogenesis. *J Comp Neurol* 2003 dec;467(1):1-10.

Buckmaster PS. Mossy Fiber Sprouting in the Dentate Gyrus. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV, editors. *Jasper's {Basic} {Mechanisms} of the {Epilepsies}*. 4th ed. Bethesda (MD); US: National Center for Biotechnology Information; 2012.

Burghardt NS, Park EH, Hen R, Fenton AA. Adult-born hippocampal neurons promote cognitive flexibility in mice. *Hippocampus* 2012 sep;22(9):1795-1808.

Butler CR, Boychuk JA, Smith BN. Effects of Rapamycin Treatment on Neurogenesis and Synaptic Reorganization in the Dentate Gyrus after Controlled Cortical Impact Injury in Mice. *Front Syst Neurosci* 2015 nov;9.

Butterbaugh G, Olejniczak P, Roques B, Costa R, Rose M, Fisch B, Carey M, Thomson J, Skinner J. Lateralization of temporal lobe epilepsy and learning disabilities, as defined by disability-related civil rights law. *Epilepsia* 2004 aug;45(8):963-970.

Bzdok D, Laird AR, Zilles K, Fox PT, Eickhoff SB. An investigation of the structural, connectional, and functional subspecialization in the human amygdala. *Hum Brain Mapp* 2013 dec;34(12):3247-3266.

Cameron HA, McKay RD. Adult neurogenesis produces a large pool of new granule cells in the dentate gyrus. *J Comp Neurol* 2001 jul;435(4):406-417.

Canpolat M, Per H, Gumus H, Yikilmaz A, Unal E, Patiroglu T, Cinar L, Kurtsoy A, Kumandas S. Rapamycin has a beneficial effect on controlling epilepsy in children with tuberous sclerosis complex: results of 7 children from a cohort of 86. *Childs Nerv Syst* 2014 feb;30(2):227-240.

Caruncho HJ, Brymer K, Romay-Tallán R, Mitchell MA, Rivera-Baltanás T, Botterill J, et al. Reelin-Related Disturbances in Depression: Implications for Translational Studies. *Front Cell Neurosci* 2016 feb;10.

Cataldi M, Avoli M, de Villers-Sidani E. Resting state networks in temporal lobe epilepsy. *Epilepsia* 2013 dec;54(12):2048-2059.

Cavazos JE, Das I, Sutula TP. Neuronal loss induced in limbic pathways by kindling: evidence for induction of hippocampal sclerosis by repeated brief seizures. *J Neurosci* 1994 may;14(5 Pt 2):3106-3121.

Cavazos JE, Golarai G, Sutula TP. Mossy fiber synaptic reorganization induced by kindling: time course of development, progression, and permanence. *J Neurosci* 1991 sep;11(9):2795-2803.

Chambers RA, Potenza MN, Hoffman RE, Miranker W. Simulated apoptosis/neurogenesis regulates learning and memory capabilities of adaptive neural networks. *Neuropsychopharmacology* 2004 apr;29(4):747-758.

Chang BS, Lowenstein DH. Epilepsy. *N Engl J Med* 2003 sep;349(13):1257-1266.

Chiaia N, Foy M, Teyler TJ. The hamster hippocampal slice. II. Neuroendocrine modulation. *Behav Neurosci* 1983;97(6):839-843.

Cho K, Lybrand ZR, Ito N, Brulet R, Tafacory F, Zhang L, et al. Aberrant hippocampal neurogenesis contributes to epilepsy and associated cognitive decline. *Nat Commun* 2015 mar;6:6606.

Chung L. A Brief Introduction to the Transduction of Neural Activity into Fos Signal. *Dev Reprod* 2015 jun;19(2):61-67.

Clark RE, Squire LR. Classical conditioning and brain systems: the role of awareness. *Science* 1998 apr;280(5360):77-81.

Cleary P, Shorvon S, Tallis R. Late-onset seizures as a predictor of subsequent stroke. *Lancet* 2004 apr;363(9416):1184-1186.

Clelland C, Choi M, Romberg C, Clemenson G, Fragniere A, Tyers P, et al. A Functional Role for Adult Hippocampal Neurogenesis in Spatial Pattern Separation. *Science* 2009 jul;325(5937):210-213.

Clugnet MC, LeDoux JE. Synaptic plasticity in fear conditioning circuits: induction of LTP in the lateral nucleus of the amygdala by stimulation of the medial geniculate body. *J Neurosci* 1990 aug;10(8):2818-2824.

Collaborators, Global Burden of Disease Study 2013. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015 aug;386(9995):743-800.

Constantine-Paton M. Effects of NMDA receptor antagonists on the developing brain. *Psychopharmacol Bull* 1994;30(4):561-565.

Cruz FC, Koya E, Guez-Barber DH, Bossert JM, Lupica CR, Shaham Y, et al. New technologies for examining the role of neuronal ensembles in drug addiction and fear. *Nature Reviews Neuroscience* 2013 nov;14(11):743-754.

Curzon P, Rustay NR, Browman KE. Cued and Contextual Fear Conditioning for Rodents. In: Buccafusco JJ, editor. *Methods of {Behavior} {Analysis} in {Neuroscience}*. 2nd ed. Boca Raton (FL): CRC Press/Taylor & Francis; 2009.

Danzer SC. Depression, stress, epilepsy and adult neurogenesis. *Exp Neurol* 2012 jan;233(1):22-32.

Danzer SC, Kotloski RJ, Walter C, Hughes M, McNamara JO. Altered morphology of hippocampal dentate granule cell presynaptic and postsynaptic terminals following conditional deletion of TrkB. *Hippocampus* 2008;18(7):668-678.

Danzer SC. Neurogenesis in Epilepsy: Better to Burn Out or Fade Away? *Epilepsy Currents* 2016 jul;16(4):268-269.

Danzer SC. Adult Neurogenesis: Opening the Gates of Troy From the Inside. *Epilepsy Curr* 2015;15(5):263-264.

Davies CH, Collingridge GL. The physiological regulation of synaptic inhibition by GABAB autoreceptors in rat hippocampus. *J Physiol (Lond)* 1993 dec;472:245-265.

Deng W, Aimone JB, Gage FH. New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nature reviews. Neuroscience* 2010 may;11(5):339-350.

Dery N, Pilgrim M, Gibala M, Gillen J, Wojtowicz JM, Macqueen G, et al. Adult hippocampal neurogenesis reduces memory interference in humans: opposing effects of aerobic exercise and depression. *Frontiers in neuroscience* 2013;7:66.

Dragunow M. Presence and induction of Fos B-like immunoreactivity in neural, but not non-neural, cells in adult rat brain. *Brain Res* 1990 nov;533(2):324-328.

Dudek FE, Sutula TP. Epileptogenesis in the dentate gyrus: a critical perspective. In: Scharfman HE, editor. Progress in {Brain} {Research}: Elsevier; 2007. p. 755-773.

Dunn KW, Kamocka MM, McDonald JH. A practical guide to evaluating colocalization in biological microscopy. *Am J Physiol , Cell Physiol* 2011 apr;300(4):723-742.

Duque A, Rakic P. Different effects of BrdU and 3H-Thymidine incorporation into DNA on cell proliferation, position and fate. *J Neurosci* 2011 oct;31(42):15205-15217.

Duvarci S, Pare D. Amygdala microcircuits controlling learned fear. *Neuron* 2014 jun;82(5):966-980.

Edwards HE, MacLusky NJ, Burnham WM. The effect of seizures and kindling on reproductive hormones in the rat. *Neurosci Biobehav Rev* 2000 sep;24(7):753-762.

Edwards HE, Burnham WM, MacLusky NJ. Testosterone and its metabolites affect afterdischarge thresholds and the development of amygdala kindled seizures. *Brain Res* 1999 aug;838(12):151-157.

Eichenbaum H. DECLARATIVE MEMORY: Insights from Cognitive Neurobiology. *Annu Rev Psychol* 1997;48(1):547-572.

El-Khayat H, Shatla H, Ali GÂ, Abdulgani M, Tomoum H, Attya H. Physical and Hormonal Profile of Male Sexual Development inÂ Epilepsy. *Epilepsia* 2003 mar;44(3):447-452.

Elst LTv, Groffmann M, Ebert D, Schulze-Bonhage A. Amygdala volume loss in patients with dysphoric disorder of epilepsy. *Epilepsy Behav* 2009 sep;16(1):105-112.

Engel J. Mesial temporal lobe epilepsy: what have we learned? *Neuroscientist* 2001 aug;7(4):340-352.

Epp JR, Silva Mera R, Köhler S, Josselyn SA, Frankland PW.. Neurogenesis-mediated forgetting minimizes proactive interference. *Nat Commun* 2016 feb;7.

Fanselow MS, Dong H. Are The Dorsal and Ventral Hippocampus functionally distinct structures? *Neuron* 2010 jan;65(1):7.

Feinstein AR. The pre-therapeutic classification of co-morbidity in chronic disease. *J Chronic Dis* 1970 dec;23(7):455-468.

Fisher RS, van Emde Boas W, Blume W, Elger C, Genton P, Lee P, et al. Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 2005 apr;46(4):470-472.

Fisher RS, Velasco AL. Electrical brain stimulation for epilepsy. *Nat Rev Neurol* 2014 may;10(5):261-270.

Forsgren L. Prevalence of epilepsy in adults in northern Sweden. *Epilepsia* 1992 jun;33(3):450-458.

Fournier NM, Andersen DR, Botterill JJ, Sterner EY, Lussier AL, Caruncho HJ, et al. The effect of amygdala kindling on hippocampal neurogenesis coincides with decreased reelin and DISC1 expression in the adult dentate gyrus. *Hippocampus* 2010 may;20(5):659-671.

Fournier NM, Botterill JJ, Marks WN, Guskjolen AJ, Kalynchuk LE. Impaired recruitment of seizure-generated neurons into functional memory networks of the adult dentate gyrus following long-term amygdala kindling. *Exp Neurol* 2013 jun;244:96-104.

Frankland PW, Köhler S, Josselyn SA.. Hippocampal neurogenesis and forgetting. *Trends Neurosci* 2013 sep;36(9):497-503.

Frasca A, Aalbers M, Frigerio F, Fiordaliso F, Salio M, Gobbi M, et al. Misplaced NMDA receptors in epileptogenesis contribute to excitotoxicity. *Neurobiol Dis* 2011 aug;43(2):507-515.

Frick KM, Kim J, Tuscher JJ, Fortress AM. Sex steroid hormones matter for learning and memory: estrogenic regulation of hippocampal function in male and female rodents. *Learn Memory* 2015 sep;22(9):472-493.

Frye CA, Reed TA. Androgenic neurosteroids: anti-seizure effects in an animal model of epilepsy. *Psychoneuroendocrinology* 1998 may;23(4):385-399.

Gage FH. Mammalian neural stem cells. *Science (New York, N.Y.)* 2000 feb;287(5457):1433-1438.

Gaitatzis A, Trimble MR, Sander JW. The psychiatric comorbidity of epilepsy. *Acta Neurol Scand* 2004 oct;110(4):207-220.

Ge S, Yang C, Hsu K, Ming G, Song H. A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron* 2007 may;54(4):559-566.

Gilbert TH, Hannesson DK, Corcoran ME. Hippocampal kindled seizures impair spatial cognition in the Morris water maze. *Epilepsy Res* 2000 feb;38(2-3):115-125.

Gilliam F, Hecimovic H, Sheline Y. Psychiatric comorbidity, health, and function in epilepsy. *Epilepsy Behav* 2003 dec;4 Suppl 4:S26-30.

Gil-Mohapel J, Simpson JM, Ghilan M, Christie BR. Neurogenesis in Huntington's disease: can studying adult neurogenesis lead to the development of new therapeutic strategies? *Brain Res* 2011 aug;1406:84-105.

Gloor P, Olivier A, Quesney LF, Andermann F, Horowitz S. The role of the limbic system in experiential phenomena of temporal lobe epilepsy. *Ann Neurol* 1982 aug;12(2):129-144.

Goddard GV, McIntyre DC, Leech CK. A permanent change in brain function resulting from daily electrical stimulation. *Exp Neurol* 1969 nov;25(3):295-330.

Goldberg EM, Coulter DA. Mechanisms of epileptogenesis: a convergence on neural circuit dysfunction. *Nat Rev Neurosci* 2013 may;14(5):337-349.

Good M, Day M, Muir JL. Cyclical changes in endogenous levels of oestrogen modulate the induction of LTD and LTP in the hippocampal CA1 region. *Eur J Neurosci* 1999;11(12):4476-4480.

Goosens KA, Maren S. Contextual and auditory fear conditioning are mediated by the lateral, basal, and central amygdaloid nuclei in rats. *Learn Memory* 2001 jun;8(3):148-155.

Gorantla VR, Pemminati S, Bond V, Meyers DG, Millis RM. Effects of Swimming Exercise on Learning and Memory in the Kainate-Lesion Model of Temporal Lobe Epilepsy. *J Clin Diagn Res* 2016 nov;10(11):F01-F05.

Gould E, Beylin A, Tanapat P, Reeves A, Shors TJ. Learning enhances adult neurogenesis in the hippocampal formation. *Nat Neurosci* 1999 mar;2(3):260-265.

Greenberg ME, Ziff EB. Stimulation of 3T3 cells induces transcription of the c-fos proto-oncogene. *Nature* 1984 oct;311(5985):433-438.

Gu Y, Arruda-Carvalho M, Wang J, Janoschka S, Josselyn S, Frankland P, et al. Optical controlling reveals time-dependent roles for adult-born dentate granule cells. *Nat Neurosci* 2012 dec;15(12):1700-1706.

Gupta A, Naorem T. Cognitive retraining in epilepsy. *Brain Inj* 2003 feb;17(2):161-174.

Guzowski JF, Setlow B, Wagner EK, McGaugh JL. Experience-dependent gene expression in the rat hippocampus after spatial learning: a comparison of the immediate-early genes Arc c-fos, and zif268. *J Neurosci* 2001 jul;21(14):5089-5098.

Hökfelt T, Ungerstedt U. Electron and fluorescence microscopical studies on the nucleus caudatus putamen of the rat after unilateral lesions of ascending nigro-neostriatal dopamine neurons. *Acta Physiol Scand* 1969 aug;76(4):415-426.

Haag A, Barth S, Zibelius M, Hermsen A, Menzler K, Oertel WH, et al. Memory for public events in patients with unilateral temporal lobe epilepsy. *Epilepsy Behav* 2010 feb;17(2):246-251.

Haag A, Barth S, Zibelius M, Hermsen A, Menzler K, Oertel WH, Hamer HM, Rosenow F, Knake S. Memory for public events in patients with unilateral temporal lobe epilepsy. *Epilepsy Behav* 2010 feb;17(2):246-251.

- Hamed SA. Neuronal plasticity: implications in epilepsy progression and management. *Drug Dev Res* 2007 dec;68(8):498-511.
- Hannesson DK, Mohapel P, Corcoran ME. Dorsal hippocampal kindling selectively impairs spatial learning/short-term memory. *Hippocampus* 2001;11(3):275-286.
- Hanson ND, Owens MJ, Nemeroff CB. Depression, antidepressants, and neurogenesis: a critical reappraisal. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 2011 dec;36(13):2589-2602.
- Hattiangady B, Shetty AK. Implications of decreased hippocampal neurogenesis in chronic temporal lobe epilepsy. *Epilepsia* 2008 jun;49:26-41.
- Hauser WA, Annegers JF, Kurland LT. Prevalence of epilepsy in Rochester Minnesota: 1940-1980. *Epilepsia* 1991 aug;32(4):429-445.
- Hayashi F, Takashima N, Murayama A, Inokuchi K. Decreased postnatal neurogenesis in the hippocampus combined with stress experience during adolescence is accompanied by an enhanced incidence of behavioral pathologies in adult mice. *Mol Brain*; 2008. p. 22.
- He X, Kotloski R, Nef S, Luikart BW, Parada LF, McNamara JO. Conditional Deletion of TrkB but Not BDNF Prevents Epileptogenesis in the Kindling Model. *Neuron* 2004 jul;43(1):31-42.
- Hellwig S, Gutmann V, Trimble MR, van Elst LT. Cerebellar volume is linked to cognitive function in temporal lobe epilepsy: a quantitative MRI study. *Epilepsy & behavior : E&B* 2013 aug;28(2):156-162.
- Henshall DC. Apoptosis signalling pathways in seizure-induced neuronal death and epilepsy. *Biochem Soc Trans* 2007 apr;35(Pt 2):421-423.
- Herzog AG, Klein P, Jacobs AR. Testosterone versus testosterone and testolactone in treating reproductive and sexual dysfunction in men with epilepsy and hypogonadism. *Neurology* 1998 mar;50(3):782-784.
- Herzog AG, Frye CA. Seizure exacerbation associated with inhibition of progesterone metabolism. *Ann Neurol* 2003 mar;53(3):390-391.
- Hlinák Z, Krejčí I. MK-801 induced amnesia for the elevated plus-maze in mice. *Behav Brain Res* 2002 apr;131(1-2):221-225.
- Hoffman GE, Smith MS, Verbalis JG. c-Fos and Related Immediate Early Gene Products as Markers of Activity in Neuroendocrine Systems. *Front Neuroendocrinol* 1993 jul;14(3):173-213.
- Holley AJ, Lugo JN. Effects of an Acute Seizure on Associative Learning and Memory. *Epilepsy&behavior : E&B* 2016 jan;54:51.

- Houser CR, Esclapez M. Vulnerability and plasticity of the GABA system in the pilocarpine model of spontaneous recurrent seizures. *Epilepsy Res* 1996 dec;26(1):207-218.
- Hsu D. The dentate gyrus as a filter or gate: a look back and a look ahead. In: Scharfman HE, editor. *Progress in {Brain} {Research}*: Elsevier; 2007. p. 601-613.
- Hu E, Mueller E, Oliviero S, Papaioannou VE, Johnson R, Spiegelman BM. Targeted disruption of the c-fos gene demonstrates c-fos-dependent and -independent pathways for gene expression stimulated by growth factors or oncogenes. *EMBO J* 1994 jul;13(13):3094-3103.
- Hughes PE, Young D, Preston KM, Yan Q, Dragunow M. Differential regulation by MK801 of immediate-early genes, brain-derived neurotrophic factor and trk receptor mRNA induced by a kindling after-discharge. *Brain Res Mol Brain Res* 1998 jan;53(1-2):138-151.
- Ikrar T, Guo N, He K, Besnard A, Levinson S, Hill A, et al. Adult neurogenesis modifies excitability of the dentate gyrus. *Front Neural Circuits* 2013;7:204.
- Iyengar SS, LaFrancois JJ, Friedman D, Drew LJ, Denny CA, Burghardt NS, et al. Suppression of Adult Neurogenesis Increases the Acute Effects of Kainic Acid. *Exp Neurol* 2015 feb;264:135-149.
- Izquierdo I. Role of NMDA receptors in memory. *Trends Pharmacol Sci* 1991 apr;12(4):128-129.
- Izquierdo I, Medina JH. Role of the amygdala, hippocampus and entorhinal cortex in memory consolidation and expression. *Braz J Med Biol Res* 1993 jun;26(6):573-589.
- Jakubs K, Nanobashvili A, Bonde S, Ekdahl CT, Kokaia Z, Kokaia M, et al. Environment Matters: Synaptic Properties of Neurons Born in the Epileptic Adult Brain Develop to Reduce Excitability. *Neuron* 2006 dec;52(6):1047-1059.
- Jasper H, Pertuiset B, Flanigin H. EEG and cortical electrograms in patients with temporal lobe seizures. *AMA Arch Neurol Psychiatry* 1951 mar;65(3):272-290.
- Jessberger S, Parent JM. Epilepsy and Adult Neurogenesis. *Cold Spring Harb Perspect Biol* 2015 nov;7(12).
- Jokeit H, Ebner A. Effects of chronic epilepsy on intellectual functions. *Prog Brain Res* 2002;135:455-463.
- Jonas P, Lisman J. Structure, function, and plasticity of hippocampal dentate gyrus microcircuits. *Front. Neural Circuits* 2014;8.
- Kalynchuk LE, Pinel JP, Treit D. Long-term kindling and interictal emotionality in rats: effect of stimulation site. *Brain Res* 1998 jan;779(1-2):149-157.

Kandel E. Principles of Neuronal Science. Fourth edition ed.; 2000.

Keezer MR, Sisodiya SM, Sander JW. Comorbidities of epilepsy: current concepts and future perspectives. *The Lancet Neurology* 2016 jan;15(1):106-115.

Kirkby RJ. Effects of lesions of the caudate nucleus or frontal neocortex on drug-induced seizures in the rat. *Psychobiology* 1977 sep;5(3):359-363.

Knoth R, Singec I, Ditter M, Pantazis G, Capetian P, Meyer RP, et al. Murine features of neurogenesis in the human hippocampus across the lifespan from 0 to 100 years. *PLoS ONE* 2010 jan;5(1):e8809.

Koek W, Woods JH, Colpaert FC. N-methyl-D-aspartate antagonism and phencyclidine-like activity: a drug discrimination analysis. *J Pharmacol Exp Ther* 1990 jun;253(3):1017-1025.

Kokaia M, Aebischer P, Elmér E, Bengzon J, Kalén P, Kokaia Z, Lindvall O. Seizure suppression in kindling epilepsy by intracerebral implants of GABA- but not by noradrenaline-releasing polymer matrices. *Exp Brain Res* 1994;100(3):385-394.

Kokaia M. Seizure-induced neurogenesis in the adult brain. *Eur J Neurosci* 2011 mar;33(6):1133-1138.

Kotloski R, Lynch M, Lauersdorf S, Sutula T. Repeated brief seizures induce progressive hippocampal neuron loss and memory deficits. *Prog Brain Res* 2002;135:95-110.

Kubik S, Miyashita T, Guzowski JF. Using immediate-early genes to map hippocampal subregional functions. *Learning & memory (Cold Spring Harbor, N.Y.)* 2007 nov;14(11):758-770.

Kuhn HG, Winkler J, Kempermann G, Thal LJ, Gage FH. Epidermal growth factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain. *J Neurosci* 1997 aug;17(15):5820-5829.

Kullmann DM. What's wrong with the amygdala in temporal lobe epilepsy? *Brain* 2011 oct;134(10):2800-2801.

La Grutta V, Sabatino M, Gravante G, Morici G, Ferraro G, La Grutta G. A study of caudate inhibition on an epileptic focus in the cat hippocampus. *Arch Int Physiol Biochim* 1988 jun;96(2):113-120.

LaBar KS, LeDoux JE, Spencer DD, Phelps EA. Impaired fear conditioning following unilateral temporal lobectomy in humans. *Journal of Neuroscience* 1995;15(10):6846-6855.

LaBar KS, Gatenby JC, Gore JC, LeDoux JE, Phelps EA. Human Amygdala Activation during Conditioned Fear Acquisition and Extinction: a Mixed-Trial fMRI Study. *Neuron* 1998 may;20(5):937-945.

Langton JM, Kim JH, Nicholas J, Richardson R. The effect of the NMDA receptor antagonist MK-801 on the acquisition and extinction of learned fear in the developing rat. *Learn Memory* 2007 oct;14(10):665-668.

LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci* 2000;23:155-184.

LeDoux JE, Cicchetti P, Xagoraris A, Romanski LM. The lateral amygdaloid nucleus: sensory interface of the amygdala in fear conditioning. *J Neurosci* 1990 apr;10(4):1062-1069.

Li F, Tsien JZ. Memory and the NMDA Receptors. *N Engl J Med* 2009 jul;361(3):302-303.

Liang KC, Lin MH, Tyan YM. Involvement of amygdala N-methyl-D-aspartate receptors in long-term retention of an inhibitory avoidance response in rats. *Chin J Physiol* 1993;36(1):47-56.

Liigant A, Haldre S, Oun A, Linnamägi U, Saar A, Asser T, Kaasik AE. Seizure disorders in patients with brain tumors. *Eur Neurol* 2001;45(1):46-51.

Lippman S, Manshadi M, Wehry M, Byrd R, Past W, Keller W, et al. 1,250 electroconvulsive treatments without evidence of brain injury. *Br J Psychiatry* 1985 aug;147:203-204.

Liu D, Bei D, Parmar H, Matus A. Activity-regulated, cytoskeleton-associated protein (Arc) is essential for visceral endoderm organization during early embryogenesis. *Mech Dev* 2000 apr;92(2):207-215.

Lledo P, Alonso M, Grubb MS. Adult neurogenesis and functional plasticity in neuronal circuits. *Nature Reviews Neuroscience* 2006 mar;7(3):179-193.

Lopez-Rojas J, Kreutz MR. Mature granule cells of the dentate gyrus are "Passive bystanders or principal performers in hippocampal function? *Neuroscience & Biobehavioral Reviews* 2016 may;64:167-174.

Lyford GL, Yamagata K, Kaufmann WE, Barnes CA, Sanders LK, Copeland NG, et al. Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron* 1995 feb;14(2):433-445.

Müller MC, Osswald M, Tinnes S, Häussler U, Jacobi A, Förster E, Frotscher M, Haas CA. Exogenous reelin prevents granule cell dispersion in experimental epilepsy. *Exp Neurol* 2009 apr;216(2):390-397.

Mahner S, Baasch C, Schwarz J, Hein S, Wölber L, Jänicke F, Milde-Langosch K. C-Fos expression is a molecular predictor of progression and survival in epithelial ovarian carcinoma. *Br J Cancer* 2008 oct;99(8):1269-1275.

Malenka RC. Synaptic plasticity in the hippocampus: LTP and LTD. *Cell* 1994 aug;78(4):535-538.

Manzanero S, Erion JR, Santro T, Steyn FJ, Chen C, Arumugam TV, et al. Intermittent fasting attenuates increases in neurogenesis after ischemia and reperfusion and improves recovery. *J Cereb Blood Flow Metab*; 2014. p. 897-905.

Maren S. Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* 2001;24:897-931.

Markwardt S, Overstreet-Wadiche L. GABAergic signalling to adult-generated neurons. *J Physiol (Lond)* 2008 aug;586(Pt 16):3745-3749.

McAuley JW, Elliott JO, Patankar S, Hart S, Long L, Moore JL, et al. Comparing patients' and practitioners' views on epilepsy concerns: a call to address memory concerns. *Epilepsy Behav* 2010 dec;19(4):580-583.

McIntyre DC. Effects of focal vs generalized kindled convulsions from anterior neocortex or amygdala on CER acquisition in rats. *Physiol Behav* 1979 nov;23(5):855-859.

Messaoudi E, Kanhema T, Soulé J, Tiron A, Dageyte G, da Silva B, Bramham CR.. Sustained Arc/Arg3.1 Synthesis Controls Long-Term Potentiation Consolidation through Regulation of Local Actin Polymerization in the Dentate Gyrus In Vivo. 2007 sep.

Milde-Langosch K. The Fos family of transcription factors and their role in tumorigenesis. *Eur J Cancer* 2005 nov;41(16):2449-2461.

Minatohara K, Akiyoshi M, Okuno H. Role of Immediate-Early Genes in Synaptic Plasticity and Neuronal Ensembles Underlying the Memory Trace. *Front Mol Neurosci* 2015;8.

Ming G, Song H. Adult Neurogenesis in the Mammalian Brain: Significant Answers and Significant Questions. *Neuron* 2011 may;70(4):687-702.

Miserendino MJD, Sananes CB, Melia KR, Davis M. Blocking of acquisition but not expression of conditioned fear-potentiated startle by NMDA antagonists in the amygdala. *Nature* 1990 jun;345(6277):716-718.

Moore PM, Baker GA. The neuropsychological and emotional consequences of living with intractable temporal lobe epilepsy: implications for clinical management. *Seizure* 2002 jun;11(4):224-230.

Morris JS, Frith CD, Perrett DI, Rowland D, Young AW, Calder AJ, et al. A differential neural response in the human amygdala to fearful and happy facial expressions. *Nature* 1996 oct;383(6603):812-815.

Moser MB, Moser EI, Forrest E, Andersen P, Morris RG. Spatial learning with a minilab in the dorsal hippocampus. *Proc Natl Acad Sci U S A* 1995 oct;92(21):9697-9701.

Narbone MC, Ruello C, Oliva A, Baviera G, D'Amico D, Bramanti P, et al. Hormonal dysregulation and catamenial epilepsy. *Funct Neurol* 1990;5(1):49-53.

- Newcomer JW, Farber NB, Olney JW. NMDA receptor function, memory, and brain aging. *Dialogues Clin Neurosci* 2000 sep;2(3):219-232.
- Obenaus A, Esclapez M, Houser CR. Loss of glutamate decarboxylase mRNA-containing neurons in the rat dentate gyrus following pilocarpine-induced seizures. *Journal of Neuroscience* 1993;13(10):4470-4485.
- O'Donnell A, Odrowaz Z, Sharrocks AD. Immediate-early gene activation by the MAPK pathways: what do and don't we know? *Biochem Soc Trans* 2012 feb;40(1):58-66.
- Oomen CA, Bekinschtein P, Kent BA, Saksida LM, Bussey TJ. Adult hippocampal neurogenesis and its role in cognition. *Wiley Interdiscip Rev Cogn Sci* 2014 sep;5(5):573-587.
- Overstreet-Wadiche LS, Bromberg DA, Bensen AL, Westbrook GL. Seizures accelerate functional integration of adult-generated granule cells. *J Neurosci* 2006 apr;26(15):4095-4103.
- Oyegbile TO, Dow C, Jones J, Bell B, Rutecki P, Sheth R, et al. The nature and course of neuropsychological morbidity in chronic temporal lobe epilepsy. *Neurology* 2004 may;62(10):1736-1742.
- Parent JM, Yu TW, Leibowitz RT, Geschwind DH, Sloviter RS, Lowenstein DH. Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J Neurosci* 1997 may;17(10):3727-3738.
- Parent JM. Adult neurogenesis in the intact and epileptic dentate gyrus. *Prog Brain Res* 2007;163:529-540.
- Parent JM. The role of seizure-induced neurogenesis in epileptogenesis and brain repair. *Epilepsy Res* 2002 jun;50(1-2):179-189.
- Parent JM, Valentin VV, Lowenstein DH. Prolonged seizures increase proliferating neuroblasts in the adult rat subventricular zone-olfactory bulb pathway. *J Neurosci* 2002 apr;22(8):3174-3188.
- Pekcec A, Potschka H. Newborn neurons with hilar basal dendrites hallmark epileptogenic networks. *Neuroreport* 2007 apr;18(6):585-589.
- Perrine K, Hermann BP, Meador KJ, Vickrey BG, Cramer JA, Hays RD, et al. The relationship of neuropsychological functioning to quality of life in epilepsy. *Arch Neurol* 1995 oct;52(10):997-1003.
- Piatti VC, Ewell LA, Leutgeb JK. Neurogenesis in the dentate gyrus: carrying the message or dictating the tone. *Front Neurosci* 2013 apr;7.
- Pierce RC, Kalivas PW. A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. *Brain Res Brain Res Rev* 1997 oct;25(2):192-216.

- Pinel JPJ, Rovner LI. Experimental epileptogenesis: Kindling-induced epilepsy in rats. *Exp Neurol* 1978 jan;58(2):190-202.
- Pitkänen A, Tuunanen J, Kälviäinen R, Partanen K, Salmenperä T.. Amygdala damage in experimental and human temporal lobe epilepsy. *Epilepsy Res* 1998 sep;32(1-2):233-253.
- Pitkänen A, Stefanacci L, Farb CR, Go G-, Ledoux JE, Amaral DG. Intrinsic connections of the rat amygdaloid complex: Projections originating in the lateral nucleus. *J Comp Neurol* 1995 may;356(2):288-310.
- Plath N, Ohana O, Dammermann B, Errington ML, Schmitz D, Gross C, et al. Arc/Arg3.1 is essential for the consolidation of synaptic plasticity and memories. *Neuron* 2006 nov;52(3):437-444.
- Pratt GD, Kokaia M, Bengzon J, Kokaia Z, Fritschy JM, Möhler H, Lindvall O. Differential regulation of N-methyl-d-aspartate receptor subunit messenger RNAs in kindling-induced epileptogenesis. *Neuroscience* 1993 nov;57(2):307-318.
- Prince HK, Conn PJ, Blackstone CD, Haganir RL, Levey AI. Down-regulation of AMPA receptor subunit GluR2 in amygdaloid kindling. *J Neurochem* 1995 jan;64(1):462-465.
- Psatta DM. Control of chronic experimental focal epilepsy by feedback caudatum stimulations. *Epilepsia* 1983 aug;24(4):444-454.
- Quartermain D, Mower J, Rafferty MF, Herting RL, Lanthorn TH. Acute but not chronic activation of the NMDA-coupled glycine receptor with D-cycloserine facilitates learning and retention. *Eur J Pharmacol* 1994 may;257(1-2):7-12.
- Rakic L, Buchwald NA, Wyers EJ. Induction of seizures by stimulation of the caudate nucleus. *Clinical Neurophysiology* 1962 dec;14(6):809-823.
- Ramírez-Amaya V, Vazdarjanova A, Mikhael D, Rosi S, Worley PF, Barnes CA.. Spatial Exploration-Induced Arc mRNA and Protein Expression: Evidence for Selective Network-Specific Reactivation. *J Neurosci* 2005 feb;25(7):1761-1768.
- Reddy DS. Testosterone modulation of seizure susceptibility is mediated by neurosteroids 3alpha-androstanediol and 17beta-estradiol. *Neuroscience* 2004;129(1):195-207.
- Reddy DS. Mass Spectrometric Assay and Physiological-Pharmacological Activity of Androgenic Neurosteroids. *Neurochem Int* 2008;52(4-5):541-553.
- Reid IC, Stewart CA. Seizures, memory and synaptic plasticity. *Seizure* 1997 oct;6(5):351-359.
- Rektor I, Kuba R, Brzdil M. Interictal and ictal EEG activity in the basal ganglia: an SEEG study in patients with temporal lobe epilepsy. *Epilepsia* 2002 mar;43(3):253-262.

Rektor I, Kuba R, Brázdil M, Chrastina J.. Do the basal ganglia inhibit seizure activity in temporal lobe epilepsy? *Epilepsy Behav* 2012 sep;25(1):56-59.

Ribak CE, Shapiro LA, Yan X, Dashtipour K, Nadler JV, Obenaus A, et al. Seizure-induced formation of basal dendrites on granule cells of the rodent dentate gyrus. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV, editors. *Jasper's {Basic} {Mechanisms} of the {Epilepsies}*. 4th ed. Bethesda (MD); US: National Center for Biotechnology Information; 2012.

Rickard NS, Poot AC, Gibbs ME, Ng KT. Both non-NMDA and NMDA glutamate receptors are necessary for memory consolidation in the day-old chick. *Behav Neural Biol* 1994 jul;62(1):33-40.

Rogers JT, Rusiana I, Trotter J, Zhao L, Donaldson E, Pak DTS, et al. Reelin supplementation enhances cognitive ability, synaptic plasticity, and dendritic spine density. *Learn Memory* 2011 sep;18(9):558-564.

Rosse RB, Mastropaolo J, Novitzki MR, Deutsch SI. Depot testosterone attenuates the anticonvulsant effect of flurazepam in mice. *Psychoneuroendocrinology* 1990;15(1):83-85.

Rowell JJ, Ragsdale CW. BrdU Birth Dating Can Produce Errors in Cell Fate Specification in Chick Brain Development. *J Histochem Cytochem* 2012 nov;60(11):801-810.

Ruxton GD, Beauchamp G. Time for some a priori thinking about post hoc testing. *Behav Ecol* 2008 may;19(3):690-693.

Santhakumar V, Aradi I, Soltesz I. Role of Mossy Fiber Sprouting and Mossy Cell Loss in Hyperexcitability: A Network Model of the Dentate Gyrus Incorporating Cell Types and Axonal Topography. *J Neurophysiol* 2005 jan;93(1):437-453.

Sayin n, Rutecki P, Sutula T. NMDA-dependent currents in granule cells of the dentate gyrus contribute to induction but not permanence of kindling. *J Neurophysiol* 1999 feb;81(2):564-574.

Scharfman HE, Sollas AL, Goodman JH. Spontaneous recurrent seizures after pilocarpine-induced status epilepticus activate calbindin-immunoreactive hilar cells of the rat dentate gyrus. *Neuroscience* 2002;111(1):71-81.

Scharfman HE, Kim M, Hintz TM, MacLusky NJ. Seizures and reproductive function: insights from female rats with epilepsy. *Ann Neurol* 2008 dec;64(6):687-697.

Scharfman HE, MacLusky NJ. Estrogen and brain-derived neurotrophic factor (BDNF) in hippocampus: complexity of steroid hormone-growth factor interactions in the adult CNS. *Front Neuroendocrinol* 2006 dec;27(4):415-435.

Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature* 2004 may;429(6988):184-187.

Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature* 2004 may;429(6988):184-187.

Schmitz B. Depression and Mania in Patients with Epilepsy. *Epilepsia* 2005 jun;46:45-49.

Schoenfeld TJ, Cameron HA. Adult Neurogenesis and Mental Illness. *Neuropsychopharmacology* 2014 sep;40(1):113-128.

Schumacher A, Vlassov E, Ito R. The ventral hippocampus, but not the dorsal hippocampus is critical for learned approach-avoidance decision making. *Hippocampus* 2016 apr;26(4):530-542.

Schwarcz R, Witter MP. Memory impairment in temporal lobe epilepsy: the role of entorhinal lesions. *Epilepsy Res* 2002 jun;50(1-2):161-177.

Seress L, Abraham H, Horváth Z, Dácsi T, Janszky J, Klemm J, et al. Survival of mossy cells of the hippocampal dentate gyrus in humans with mesial temporal lobe epilepsy. *J Neurosurg* 2009 dec;111(6):1237-1247.

Shin C, McNamara JO, Morgan JI, Curran T, Cohen DR. Induction of c-fos mRNA expression by afterdischarge in the hippocampus of naive and kindled rats. *J Neurochem* 1990 sep;55(3):1050-1055.

Shirvalkar P, Seth M, Schiff ND, Herrera DG. Cognitive enhancement with central thalamic electrical stimulation. *Proc Natl Acad Sci U S A* 2006 nov;103(45):17007-17012.

Sloviter RS. Decreased hippocampal inhibition and a selective loss of interneurons in experimental epilepsy. *Science* 1987;235(4784):73-76.

Sloviter RS, Zappone CA, Harvey BD, Bumanglag AV, Bender RA, Frotscher M. Dormant basket cell hypothesis revisited: Relative vulnerabilities of dentate gyrus mossy cells and inhibitory interneurons after hippocampal status epilepticus in the rat. *J Comp Neurol* 2003;459(1):44-76.

Smith MD, Jones LS, Wilson MA. Sex differences in hippocampal slice excitability: role of testosterone. *Neuroscience* 2002 feb;109(3):517-530.

Snyder JS, Hong NS, McDonald RJ, Wojtowicz JM. A role for adult neurogenesis in spatial long-term memory. *Neuroscience* 2005;130(4):843-852.

Spalding KL, Bergmann O, Alkass K, Bernard S, Salehpour M, Huttner HB, et al. Dynamics of hippocampal neurogenesis in adult humans. *Cell* 2013 jun;153(6):1219-1227.

Spampanato J, Polepalli J, Sah P. Interneurons in the basolateral amygdala. *Neuropharmacology* 2011 apr;60(5):765-773.

- Spehlmann R, Norcross K, Grimmer EJ. GABA in the caudate nucleus: A possible synaptic transmitter of interneurons. *Experientia* 1977 may;33(5):623-625.
- Sramka M, Fritz G, Galanda M, Nájdvorník P. Some observations in treatment stimulation of epilepsy. *Acta Neurochir* 1976(23 Suppl):257-262.
- Steward O, Farris S, Pirbhoy PS, Darnell J, Driesche SJV. Localization and local translation of Arc/Arg3.1 mRNA at synapses: some observations and paradoxes. *Front Mol Neurosci* 2014;7.
- Irene Suárez-Pereira and Ángel M Carrióna.. Updating stored memory requires adult hippocampal neurogenesis. *Scientific Reports*, Published online: 11 September 2015;
- Sutula T, Koch J, Golarai G, Watanabe Y, McNamara JO. NMDA receptor dependence of kindling and mossy fiber sprouting: evidence that the NMDA receptor regulates patterning of hippocampal circuits in the adult brain. *J Neurosci* 1996 nov;16(22):7398-7406.
- Sutula TP. Mechanisms of epilepsy progression: current theories and perspectives from neuroplasticity in adulthood and development. *Epilepsy Res* 2004 aug;60(2-3):161-171.
- Swanson LW, Cowan WM. An autoradiographic study of the organization of the efferent connections of the hippocampal formation in the rat. *J Comp Neurol* 1977 mar;172(1):49-84.
- Teixeira CM, Kron MM, Masachs N, Zhang H, Lagace DC, Martinez A, et al. Cell-autonomous inactivation of the Reelin pathway impairs adult neurogenesis in the hippocampus. *J Neurosci* 2012 aug;32(35):12051-12065.
- Thom M. Review: Hippocampal sclerosis in epilepsy: a neuropathology review. *Neuropathol Appl Neurobiol* 2014 aug;40(5):520-543.
- Toothaker LE. *Multiple Comparison Procedures*. : SAGE; 1993.
- Tu Y, Huang D, Shiah S, Wang J, Lin W. Regulation of c-Fos Gene Expression by NF- $\kappa$ B: A p65 Homodimer Binding Site in Mouse Embryonic Fibroblasts but Not Human HEK293 Cells. *PLOS ONE* 2013 dec;8(12):e84062.
- Tulchinsky E. Fos family members: regulation, structure and role in oncogenic transformation. *Histol Histopathol* 2000 jul;15(3):921-928.
- Turski L, Cavalheiro EA, Bortolotto ZA, Ikonomidou-Turski C, Kleinrok Z, Turski WA. Dopamine-sensitive anticonvulsant site in the rat striatum. *J Neurosci* 1988 nov;8(11):4027-4037.
- Turski L, Meldrum BS, Cavalheiro EA, Calderazzo-Filho LS, Bortolotto ZA, Ikonomidou-Turski C, et al. Paradoxical anticonvulsant activity of the excitatory amino acid N-methyl-D-aspartate in the rat caudate-putamen. *Proc Natl Acad Sci U S A* 1987 mar;84(6):1689-1693.

Van Strien NM, Cappaert NLM, Witter MP. The anatomy of memory: an interactive overview of the parahippocampal-hippocampal network. *Nat Rev Neurosci* 2009 apr;10(4):272-282.

Vazdarjanova A, McGaugh JL. Basolateral amygdala is not critical for cognitive memory of contextual fear conditioning. *Proc Natl Acad Sci U S A* 1998 dec;95(25):15003-15007.

Vella N, Ferraro G, Caravaglios G, Aloisio A, Sabatino M, La Grutta V. A feature of caudate control of focal hippocampal epilepsy: evidence for an anterograde pathway. *Exp Brain Res* 1991;85(1):240-242.

Verrotti A, Latini G, Manco R, Simone MD, Chiarelli F. Influence of sex hormones on brain excitability and epilepsy. *J Endocrinol Invest* 2007 oct;30(9):797-803.

Victor Nadler J. Kainic acid as a tool for the study of temporal lobe epilepsy. *Life Sci* 1981 nov;29(20):2031-2042.

Wanisch K, Tang J, Mederer A, Wotjak CT. Trace fear conditioning depends on NMDA receptor activation and protein synthesis within the dorsal hippocampus of mice. *Behav Brain Res* 2005 feb;157(1):63-69.

Warren SG, Humphreys AG, Juraska JM, Greenough WT. LTP varies across the estrous cycle: enhanced synaptic plasticity in proestrus rats. *Brain Res* 1995 dec;703(1-2):26-30.

Wiskott L, Rasch MJ, Kempermann G. A functional hypothesis for adult hippocampal neurogenesis: avoidance of catastrophic interference in the dentate gyrus. *Hippocampus* 2006;16(3):329-343.

Won SJ, Kim SH, Xie L, Wang Y, Mao XO, Jin K, et al. Reelin-deficient mice show impaired neurogenesis and increased stroke size. *Exp Neurol* 2006 mar;198(1):250-259.

Wong M. Rapamycin for Treatment of Epilepsy: Antiseizure Antiepileptogenic Both or Neither? *Epilepsy Curr* 2011;11(2):66-68.

Woolley CS. Estradiol facilitates kainic acid-induced, but not flurothyl-induced, behavioral seizure activity in adult female rats. *Epilepsia* 2000 may;41(5):510-515.

Zhang J, Jiao J. Molecular Biomarkers for Embryonic and Adult Neural Stem Cell and Neurogenesis. *Biomed Res Int* 2015;2015.

Zhang J, Zhang D, McQuade JS, Behbehani M, Tsien JZ, Xu M. c-fos regulates neuronal excitability and survival. *Nat Genet* 2002 apr;30(4):416-420.

Zhao C, Deng W, Gage FH. Mechanisms and Functional Implications of Adult Neurogenesis. *Cell* 2008 feb;132(4):645-660.

Zhong Q, Ren B, Tang F. Neurogenesis in the Hippocampus of Patients with Temporal Lobe Epilepsy. *Curr Neurol Neurosci Rep* 2016 feb;16(2):20.