

---

---

# Methane and nitrous oxide emission from rice field under long-term fertilization

Qin Xiaobo<sup>1,2</sup>, Li Yu'e<sup>1,2</sup>, Wan Yunfan<sup>1,2</sup>, Gao Qingzhu<sup>1,2</sup>, Wang Hong<sup>3</sup> and Kelsey Brandt<sup>3</sup>

<sup>1</sup>Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of Agricultural Sciences, 100081 Beijing China (e-mail: [chinayrh@gmail.com](mailto:chinayrh@gmail.com))

<sup>2</sup>The key laboratory for Agro-Environment and Climate Change, Ministry of Agriculture, 100081 Beijing China

<sup>3</sup>Semiarid Prairie Agricultural Research Centre, AAFC Swift Current SK S9H3X2 Canada

---

---

## Introduction

Rice is an important crop in China, both as major food source, and an export commodity. There are many factors which affect CH<sub>4</sub> and N<sub>2</sub>O emission from rice fields. A rice experiment in the Hunan province of China with two seeding dates and 4 fertilizer treatments began in 1980. The fertilizer treatments consisted of no fertilizer, inorganic fertilizer only, or a combination of inorganic fertilizer and an organic amendment, either rice straw or hog manure. The purpose of this study is to determine the effect of these fertilizer treatments on greenhouse gas (GHG) emissions.

## Materials and methods

Four long-term fertilization treatments were selected from 2004 to observe GHG emissions. They are 1) NPKS (NPK + straw); 2) CK (no fertilizer); 3) NPK; and 4) NKM (NK + hog manure). The fertilizers applied were Urea (N), P<sub>2</sub>O<sub>5</sub> (P), K<sub>2</sub>O (K). The early rice treatment received 75 kg ha<sup>-1</sup> N initially and was top dressed with an additional 75 kg ha<sup>-1</sup> N. The late rice received 90 kg ha<sup>-1</sup> N and was top dressed with another 90 kg ha<sup>-1</sup> N. Rates of P and K were applied at 45 kg ha<sup>-1</sup> and 120 kg ha<sup>-1</sup> respectively. All fertilizers are applied as kg ha<sup>-1</sup> of product. Decomposed hog manure from a biogas container (Marsh gas-methane) which is used to collect methane gas from the fermentation process was applied at a rate of 15000 kg ha<sup>-1</sup> to the NKM treatment. The NPKS treatment received rice straw at 2625 kg ha<sup>-1</sup>.

There is one plot for each treatment, and three repetitions in one plot. We installed one sampling chamber for each repetition. In the entire experimental period from the sowing of early rice to the harvest of late rice, GHG sample were taken manually once every three days, usually 9~10 am. Concentration of GHG was analyzed by Agilent 6890N GC.

The flux of GHG was calculated by the following formula:

$$F = \rho \cdot h \cdot dC/dt \cdot 273 / (273 + T)$$

Where:

F - GHG Flux (CH<sub>4</sub>:mg·m<sup>-2</sup>·h<sup>-1</sup>, N<sub>2</sub>O:μg·m<sup>-2</sup>·h<sup>-1</sup>),

$\rho$ - Density of GHG in standard conditions  
(CH<sub>4</sub>: 0.717kg·m<sup>-3</sup>, N<sub>2</sub>O: 1.97kg·m<sup>-3</sup>)  
 $h$  - Height of chamber (m),  
 $dC/dt$  - Rate of GHG change in chamber,  
 $T$  - Average chamber temperature during sampling.

## **Results and discussion**

### **Seasonal variation of CH<sub>4</sub> flux from rice field**

There is only one big CH<sub>4</sub> flux peak during the entire rice growing season (Fig. 1). Most of CH<sub>4</sub> fluxes were emitted before drainage, after that, there was only a little CH<sub>4</sub> emission.

### **Seasonal variation of N<sub>2</sub>O flux from rice field**

The seasonal variation of N<sub>2</sub>O flux from rice field was distinct with CH<sub>4</sub> (Fig. 2). There were several N<sub>2</sub>O flux peaks in the growing season, but the largest one occurs after drainage.

### **Effect of long-term fertilization on GHG emission from rice field**

The rank of CH<sub>4</sub> average flux in different fertilization treatments is: NPKS>NKM>CK>NPK (Fig. 3). The rank of N<sub>2</sub>O average flux in different treatments from early rice is: NPK>NKM>NPKS>CK, but from late rice is: NPKS>NPK>NKM>CK (Fig. 4).

### **Relationship between CH<sub>4</sub> flux and soil temperature**

In the range of 15~30°C, CH<sub>4</sub> flux from rice field increased with the soil temperature at 10cm.

In the same temperature range, N<sub>2</sub>O flux from late rice field after drainage increasing with the soil temperature in 10cm (Fig. 5).

### **Relationship between CH<sub>4</sub> flux and soil pH**

In the range of pH5-5.8 and pH6.2-6.8, CH<sub>4</sub> flux from rice field increasing with the soil pH, there is a linear relationship between CH<sub>4</sub> flux and soil pH (Fig. 6). In the range of pH6.2-6.8, N<sub>2</sub>O flux from late rice field before drainage is increasing with the soil pH.

## **References**

Zheng X H, Wang M X, Wang Y S et al.. Comparison of manual and automatic methods for measurement of methane emission from rice paddy fields [J]. Advances in Atmospheric Sciences, 1998,15(4):569-579.

## **Acknowledgements**

The authors of this paper would like to thank Mr. Li Yuehui, Mrs. Chang Fuyuan, Yang Ruihua for their assistance with the GHGs sampling, Liao Yulin, Shi Feng, for their assistance with data collection.

Figure 1 Seasonal variation of CH<sub>4</sub> flux from rice field




Figure 2 Seasonal variation of N<sub>2</sub>O flux from rice field




Figure 3. CH<sub>4</sub> flux from every long-term fertilization treatment




Figure 4. N<sub>2</sub>O flux from every long-term fertilization treatment




Figure 5. Relationship between CH<sub>4</sub> flux and soil temperature in 10cm depth




Figure 6. Relationship between CH<sub>4</sub> flux and soil pH

