

# **The Effects of Imposing Tillage on Long-term No-till Soils**

C.D. Baan<sup>1\*</sup>, M.C.J. Grevers<sup>1</sup> & J.J. Schoenau<sup>1</sup>

<sup>1</sup>Department of Soil Science, University of Saskatchewan, Saskatoon, SK, Canada;

## **Introduction**

During the last two decades technology has advanced enough for producers to manage their soil with little or no tillage. Producers that have adopted the no till system have realized reduced erosion, increased moisture conservation, better soil structure and higher net returns (Singh and Singh, 1994; Larney *et al.*, 1997).

With maturing no-till fields, some issues have been brought forward:

- Concerns with nutrient stratification near the soil surface (Grant & Bailey, 1994)
- Perennial/herbicide resistant weeds (Derksen *et al.*, 1993)
- Low soil temperatures in the spring (Wolf and Eckert, 1999)

This may lead a producer to consider imposing tillage to alleviate one or all of these problems.

**The objective of this project was to study the effect of imposing one cycle of tillage of different intensities on a long-term no-till soil (10+years) in the Brown, Black, and Gray soil zones.**

We hypothesized that one cycle of tillage will:

- 1) encourage organic matter decomposition & nutrient release
- 2) decrease nutrient stratification, alter soil density, structure and moisture content of the surface soil
- 3) Reduce heavy thatch on soil surface and increase spring soil temperatures.

## **Materials & Methods**

### *Study Sites:*

Each site had a completely randomized design with four replications and a plot size of 24X50 feet.

Treatments included:

1. No-till: control
2. Min-till: Spring pass with 12" sweeps
3. Conv.-till: Fall + spring passes with 12" sweeps
4. Max-till: Fall + spring passes with 12" sweeps + one pass with a tandem disc.

### ***Sampling techniques:***

After tillage treatments were imposed and prior to seeding, polyvinyl chloride (PVC) cores were punched 15 cm into the soil and intact soil cores were removed. Surface (5cm depth) temperatures were also taken at this time

A punch truck was used to determine the moisture regime in the soil down to 90 cm

Fall harvest samples were analyzed for yield and nutrient content.

Following harvest another set of intact soil cores were taken

### ***Core soil analysis***

The intact soil cores collected in the spring were then subjected to a two week incubation to determine CO<sub>2</sub> production as a result of microbial respiration and decomposition. Gas samples were taken from the PVC cores which were capped with an airtight seal. Seven days into the incubation, the cores were rewetted to simulate a light rainfall.

Also during the incubation, nutrient ion supply rates were determined by plant root simulator (PRS<sup>TM</sup>) probes which use resin membranes as an ion sink (Qian and Schoenau, 2002).

After the incubation the cores were sectioned into 5cm increments, bulked, and analyzed for bulk density, pH, EC, organic matter and extractable nutrients.

Extractable P and K were determined by modified Kelowna solution and ammonium and nitrate extracted with 2M KCl.

Grain and straw yield were measured and N and P concentrations were assessed by wet digestion(Carter, 1993).

## SOIL ZONES OF SASKATCHEWAN




Figure 2. Plot sites in Brown, Black, and Gray soil zones



Figure 1. Tilling Brown soil zone site



**Figure 4. Tilled Black soil zone site**



**Figure 3. Tilled Gray soil zone site.**

## Results and Discussion


All three sites showed that the soil that was not tilled produced the least amount of CO<sub>2</sub>. Tillage exposes organic matter to attack from microorganisms and thus produces higher fluxes of CO<sub>2</sub> due to increased microbial activity and decomposition rates.

At the Central Butte site, one tillage operation in the spring appeared to reduce the supply rate of nitrate while more intensive tillage had no effect. At Rosthern and Tisdale sites, intensive tillage significantly reduced the supply rates of soil nitrate.

The reduced nitrate supply rates at Rosthern and Tisdale are most likely due to the fresh straw (high C:N ratio) incorporated during tillage. The Central Butte site had been in



**Figure 6. Incubation chambers containing intact soil cores with PRS probes inserted.**




**Figure 5. CO2 chambers with intact core sealed in and ready for gas sampling**

chem-fallow the previous year and had lower amounts of more highly weathered, degraded straw.


Shortly after tillage, soil density was reduced and increased soil temperatures of about 0.5 to 1.5 degrees C were observed. It generally only took one tillage operation (spring only) to produce these effects.

Tillage showed no significant effect on pH, E.C., and total soil organic matter.


No significant differences were noted in the grain and straw yields among the tillage treatments but there was a trend to decreased yields at Rosthern and Tisdale, possibly reflecting the N immobilization. Mixing of soil reduced nutrient stratification near the soil surface.



**Figure 7. Mean NO<sub>3</sub> in Brown Soil Zone +/- SE**



**Figure 8. Mean NO<sub>3</sub> in Black Soil Zone +/- SE**



**Figure 9. Mean NO<sub>3</sub> in Gray Soil Zone +/- 1 SE**

## Conclusions

- ❖ Soil incubation of intact cores revealed higher CO<sub>2</sub> fluxes in tilled soils than no-till soils.
- ❖ An apparent effect of tillage on enhancing immobilization of available N due to fresh straw incorporation was observed
- ❖ Soil density was reduced and temperatures were increased by tillage
- ❖ Total soil organic carbon, pH, and E.C. were unaffected by tillage
- ❖ Tillage increased uniformity of nutrient distribution with depth.

## Acknowledgements

We thank the Agricultural Development Fund (ADF) for the funding to carry out this project.

❖ **For more information contact Chris at [cdb661@mail.usask.ca](mailto:cdb661@mail.usask.ca)**

## References

- Carter, M.R. (ed) 1993. Soil Sampling and Methods of Analysis. Can. Soc. of Soil Sci. Lewis Publishers, Boca Raton, FL.
- Derksen, D.A., G.P. Lafond, A. G. Thomas, H. A. Loeppky, and C. J. Swanton. 1993. Impact of Agronomic Practices on Weed Communities: Tillage Systems. *Weed Sci.* 41:409-417.
- Grant, C.A., and L.D. Bailey. 1994. The Effect of Tillage and KCl Addition on pH, Conductance, NO<sub>3</sub>-N, P, K and Cl Distribution in the Soil Profile. *Can. J. Soil Sci.*
- Larney, F.J., E. Bremer, H.H. Janzen, A.M. Johnston, and C.W. Lindwall. 1997. Changes in Total, Mineralizable and Light Fraction Soil Organic Matter with Cropping and Tillage Intensities in Semiarid Southern Alberta, Canada. *Soil and Tillage Research.* 42:229-240.
- Qian, P., and J.J. Schoenau. 2002. Practical Applications of Ion Exchange Resins in Agricultural and Environmental Soil Research. *Can. J. Soil Sci.* 82:9-21.
- Schoenau, J. J., and C. A. Campbell. 1996. Impact of Crop Residues on Nutrient Availability in Conservation Tillage Systems. *Can. J. Plant Sci.* 76:621-626.
- Singh, H., and K.P. Singh. 1994. Nitrogen and Phosphorus Availability and Mineralization in Dryland Reduced Tillage Cultivation: Effects of Residue Placement and Chemical Fertilizer. *Soil Biol. Biochem.* 26:695-702.
- Wolfe, A.M., and D.J. Eckert. 1999. Crop Sequence and Surface Residue Effects on the Performance of No-Till Corn Grown on a Poorly Drained Soil. *Agron. J.* 91:363-367.