NITROGEN FERTILIZATION OF WHEAT SEEDED ON SUMMERFALLOW LAND

E. H. Halstead - Department of Soil Science University of Saskatchewan

The fields used in this study were selected from summerfallow fields tested in the fall of 1966 which had lower than normal nitrate-nitrogen tests. Rates of fertilizer applied were based on soil test data.

RESULTS

YIELD INCREASE TO TREATMENT (Bushels/acre)

Farmer	Soil	Check Yield	Yield Increase**			
	Туре		23-23-0	11-48-0	11-48-0	18-46-0
		Conference and the second s	and the second		33.5-0-0	Name dalam - a da ana mandra da ana ana ang sa
*Evans 1	BrVL	19.23	+ 8.31	+ 5.29	+ 6.41 .	+ 5.82
Evans 2	WL	18.17	+ 8.30	+ 6.27	+ 4.39	
*Drew	RHvC	23.70	+ 0.48	- 2.29	+ 4.28	+ 3.66
Rennick 1	RHvC	13.00	+ 8.73	- 0.48	+ 4.93	
Grandfield	LcVL	8.75	- 2.91	+ 2.78	- 0.20	
*Lazaroff	KSiL	23.99	+ 5.36	+ 2.87	+ 6.57	+10.30
Cadrin 1	HmL	33.39	+15.50	+16.25	+11.54	
Cadrin 2	HmL	52.42	+ 1.67	+ 0.08	+ 6.42	
*Halstead	AFL	25.60	+ 7.36	+ 3.56	+ 4.72	+ 4.09
*Konschuh	BrVL	29.31	+ 5.54	+ 5.63	+ 5.27	+ 5.46
Ewert	WL	20.18	+ 3.22	+ 0.34	+ 4.07	
Wilkinson	ECL	29.72	+ 4.99	+ 5.78	+12.25	
AVERAGE (12)		24.79	+ 5.55	+ 3.84	+ 5.89	
*AVERAGE (5))	24.37	+ 5.41	+ 3.01	+ 5.45	+ 5.87
						이번 물리는 것은 비가 문제했다.

** 23-23-0 applied at 80 lb/acre. Rates of N and/or P₂O₅ for other treatments as recommended by the soil test for stubble seeded crops.

SUMMARY AND RECOMMENDATIONS

In most cases, higher wheat yields were obtained where additional nitrogen above that in 11-48-0 was applied. On the average 23-23-0 or the split application of 11-48-0 + 33.5-0-0 yielded about 2 bushels/acre more than the 11-48-0 treatment. On the basis of 5 trials, 18-46-0 at rates recommended by the phosphorus test was superior to 11-48-0. The average yield for 18-46-0 was similar to the higher nitrogen applications.

Based on these results it is recommended that on summerfallow fields, with 0-24 inch nitrate-nitrogen levels of less than 45 lb/acre, that higher nitrogen ammonium phosphates such as 23-23-0, 18-46-0, 17-34-0, etc., be considered as alternatives to 11-48-0. Split application of N plus P_2O_5 should be considered for fields with very low and low soil test levels.