

---

# Nitrous Oxide Emissions from Soil Profiles Seeded with Pulse Crops

Z. Z. Zhong<sup>1</sup>, L. M. Nelson<sup>2</sup>, R. L. Lemke<sup>3</sup>

<sup>1</sup>Department of Soil Science, University of Saskatchewan. Saskatoon, SK S7N 5A8

<sup>2</sup>Department of Applied Microbiology and Food Science, University of Saskatchewan, Saskatoon, SK S7N 5A8

<sup>3</sup>Agriculture and Agri-Food Canada, Semiarid Prairie Agriculture Research Centre, Swift Current, SK S9H 3X2

---

**Key Words:** nitrous oxide emission, pulse crop, denitrification, nitrogen fixation

## Abstract

Inoculation of legumes with *Rhizobium* spp. is a common worldwide agricultural practice that is used to increase crop yield and to improve soil fertility without adding nitrogen (N) fertilizers. There is concern that N<sub>2</sub> fixation by legumes enhances emissions of nitrous oxide (N<sub>2</sub>O) – a powerful greenhouse gas (GHG). The objectives of this experiment were: 1) to measure N<sub>2</sub>O emissions from soil profiles under inoculated and non-inoculated pulse crops; 2) to investigate the relationship between N<sub>2</sub>O emissions and N<sub>2</sub> fixation by pulse crops. The experiment was carried out in the Phytotron of the University of Saskatchewan with soil flats seeded with inoculated and non-inoculated lentils and peas and fertilized spring wheat. N<sub>2</sub>O emissions from soil profiles were measured using Profile Gas Samplers (PGS), and were analyzed with a GC. Results indicated that inoculated lentils and peas did not substantially increase N<sub>2</sub>O emissions, and N<sub>2</sub>O emissions may not be associated with N<sub>2</sub> fixation under the conditions in this experiment.

## 1 Introduction

Rhizobia can invade and form nodules on the roots of leguminous plants to fix nitrogen from the atmosphere. Inoculation of legumes with *Rhizobium* spp. is a common worldwide agricultural practice used to increase crop yield and to improve soil fertility without adding nitrogen fertilizers (Bohlool et al., 1992; Peoples and Craswell, 1992). Substantial acreages of pulse crops such as field pea and lentil are grown in Western Canada. Recently, questions have arisen about whether the cultivation of pulse crops enhances N loss in the form of N<sub>2</sub>O – a powerful greenhouse gas (Svensson et al., 1991; Klemmedtsson, et al., 1991; Aulakh et al., 1991; Arrese-Igor and Aparicio-Tejo, 1992; Kilian and Werner, 1996). The enhanced N<sub>2</sub>O emissions from agricultural and natural ecosystems are believed to be caused by increasing soil N availability driven by increased fertilizer use, agricultural nitrogen (N<sub>2</sub>) fixation, and N deposition (IPCC, 2001). The current methodology for estimating N<sub>2</sub>O emissions on a national level considers N<sub>2</sub> fixation by legumes an important source of N<sub>2</sub>O emissions (IPCC, 2001). However, the contribution of inoculated pulse crops to N<sub>2</sub>O emissions under dryland conditions in

western Canadian is not well understood. The objectives of this experiment were: 1) to measure N<sub>2</sub>O emissions from soil profiles seeded with inoculated and non-inoculated lentils and peas; 2) to investigate the relationship between N<sub>2</sub>O emissions and N<sub>2</sub> fixation by lentils and field peas.

## **2 Materials and Methods**

### **2.1 Soil and soil flats**

A Chernozemic soil was collected from the Dark Brown soil zone and was homogeneously mixed with sand. The resultant soil has a texture of sandy loam with a bulk density of 1.35 g cm<sup>-3</sup>, and total C, N and P contents of 34.4, 2.44, 0.68 g kg<sup>-1</sup> soil, respectively. Using this soil, 16 soil flats (45 cm × 50 cm × 60 cm) were built for plant growth.

### **2.2 Inoculant preparation**

Rhizobial inoculants were prepared by growing two *R. leguminosarum biovar viciae* isolates 99A1 and RGP2 in yeast extract-mannitol (YEM) medium. Cultures were placed on a G10 Gyrotory Shaker (at 22±C, 150 rpm) for 3 d prior to inoculation.

### **2.3 Plant inoculation, fertilization and growth conditions**

Lentil (*Lens esculenta* Moench cv. *Milestone*) and pea (*Pisum sativum* L. cv. *Mozart*) were inoculated at a rate of 3 ml g<sup>-1</sup> of 99A1 and RGP2, respectively, at sowing date. Non-nodulated pulse crops were fertilized with NH<sub>4</sub>NO<sub>3</sub> (100 kg N ha<sup>-1</sup>). Spring wheat (*AC Barrie*), used as a reference crop, was fertilized at a rate of 100 kg N ha<sup>-1</sup>. P and K were applied at a rate of 15 kg P ha<sup>-1</sup> and 35 kg K ha<sup>-1</sup>, respectively. Each treatment was replicated 3 times.

Plants were grown in the phytotron of University of Saskatchewan (daytime: 22±C for 16 h, nighttime: 16±C for 8 h). Soil moisture was maintained at 60-80% of field capacity by monitoring with a time domain reflectometry (TDR) (Topp et al., 1996). Plants were sown on July 15, 2003. They emerged on July 22, and were harvested on October 23, 2003.

### **2.4 N<sub>2</sub>O measurement**

A multi-level Profile Gas Sampler (PGS) was constructed for gas sampling from soil profiles at different depths. Gas samples were taken once a week using a syringe and injected into evacuated tubes, and their N<sub>2</sub>O concentrations were analyzed using a gas chromatograph (GC) (Varian CP3800). The GC consists of three electron capture detectors (ECD). The carrier gas was 95% argon and 5% methane, and make-up gas was helium. The temperatures for detector, oven and injector were 370±C, 35±C, and 100±C, respectively. The software for the integrator/recorder was Varian Star 5.5. During the GC operation, a gas sample of 300 vL was introduced into the injection system for the analysis of N<sub>2</sub>O concentrations.

## 2.5 Other measurements

Nitrogen fixation by lentil and pea, as indicated by nitrogenase activities of nodules, was determined using the  $C_2H_2$  reduction assay (ARA) (Hardy et al., 1968). Plants were carefully uprooted from the soil flats so that nodules were least removed and then were placed as quickly as possible in screw-cap jars (550 ml in volume) fitted with a rubber seal for gas injections and sampling. Then, 25 ml of air was drawn out of each sealed jar and replaced with the same volume of acetylene. After 30 min of incubation (under  $25\pm C$ ), about 0.5 ml gas sample was drawn from the jar using a syringe and was injected into a gas chromatography (Model 5790A Series, Hewlett-Packard) for  $C_2H_2$  reduction analysis. Within the GC system, the gas carrier was dinitrogen gas ( $N_2$ ) with a flow rate of  $30\text{ ml min}^{-1}$ . The temperatures for oven, injector and detector were  $40\pm C$ ,  $50\pm C$  and  $150\pm C$ , respectively. The pressures for  $N_2$  and  $H_2$  and air were 160 kPa, 200 kPa, and 150 kPa, respectively.

Plant biomass was measured at different growing stages by randomly removing and drying (at  $60\pm C$  for 48 h in oven) one plant from each soil box. Plant grain yields were measured at harvest time.

## 3 Results and Discussion

### 3.1 Plant growth and nitrogenase activities

Inoculation of *R. leguminosarum* 99A1 onto lentil and RGP2 onto field pea provided N for plant growth. Nitrogen fixation by nodules, as indicated by nitrogenase activities, was significantly higher in inoculated lentil and pea than in non-inoculated plants (Table 1). However, plant biomass and grain yields were generally not significantly different between inoculated and fertilized plants.

**Table 1 Plant Growth and Nitrogenase Activity of Nodules**

Plant	Inoculant /fertilizer	Shoot biomass (g plant <sup>-1</sup> )	Root biomass (g plant <sup>-1</sup> )	Yield (g m <sup>-2</sup> )	Nitrogenase activity (vmol g <sup>-1</sup> h <sup>-1</sup> )
Lentil	99A1	5.12 " 1.59ab	0.07 " 0.04a	254.36 " 11.04b	14.06 " 1.41a
Lentil	100 kg N ha <sup>-1</sup>	2.55 " 1.37b	0.07 " 0.01a	250.10 " 24.91b	4.29 " 1.36b
Pea	RGP2	6.60 " 0.93a	0.41 " 0.44a	298.59 " 25.85a	19.21 " 5.83a
Pea	100 kg N ha <sup>-1</sup>	5.31 " 1.51a	0.09 " 0.03a	328.93 " 5.82a	2.25 " 0.38b

Nitrogenase activity and plant biomass were measured at the mid-stage of plant growth.

Values with different letter are significantly different ( $p<0.05$ ).


### 3.2 $N_2O$ production from soil profiles

Low  $N_2O$  production (0.3 -3.5 ppm) was observed 0-25 d after planting, after which  $N_2O$  concentrations leveled off and remained at the same level as the ambient atmosphere (Figs. 1-5).

Most of the N<sub>2</sub>O production accumulated in the deeper layers (15-30 cm). This may have resulted from reduced gas exchange between the deeper layers and the surface and reduced aeration in the deeper layers, favoring denitrification.

The N<sub>2</sub>O emissions from soil profiles showed the same patterns both for pulse crops and the cereal crop, and for inoculated and non-inoculated pea and lentil. Rhizobial strains 99A1 and RGP2 were not able to denitrify in pure cultures (Zhong et al., 2003). This indicated that inoculation of lentil and pea with these rhizobial strains did not enhance denitrification under the conditions of this experiment.

Nitrogen fixation by lentil and pea, as expressed by nitrogenase activities of nodules, was significantly increased in inoculated compared to non-inoculated legumes (Table 1). However, N<sub>2</sub>O production from soil profiles was not significantly different between inoculated and non-inoculated lentil and pea (Figs. 1-4). In addition, a comparison of N<sub>2</sub>O production from soil profiles between pulse crops (lentil and pea) and a cereal crop (spring wheat) indicated that the presence of pulse crops did not have an obvious impact on N<sub>2</sub>O emissions from soil profiles (Figs. 1-5). These results suggest that N<sub>2</sub>O emissions may not be associated with nitrogen fixation by pulse crops.



**Fig. 1 N<sub>2</sub>O concentrations in soil profiles of inoculated lentil**




Fig. 2 N<sub>2</sub>O concentrations in soil profiles of non-inoculated lentil




Fig. 3 N<sub>2</sub>O concentrations in soil profiles of inoculated pea




Fig. 4 N<sub>2</sub>O concentrations in soil profiles of non-inoculated pea




Fig. 5 N<sub>2</sub>O concentrations in soil profiles of fertilized wheat

## 4 Conclusion

Inoculation of *R. leguminosarum* strains 99A1 and RGP2 onto lentil and pea stimulated plant growth, but N<sub>2</sub>O production was not enhanced compared with fertilization with NH<sub>4</sub>NO<sub>3</sub>. Inoculating pulse crops with these two rhizobial strains increased soil fertility and crop yields, but did not increase N<sub>2</sub>O production. This suggests that N<sub>2</sub>O emissions may not be associated with nitrogen fixation by pulse crops under these conditions.

## References

- Arrese-Igor, C., Aparicio-Tejo, P.M., 1992. Denitrification and respiration in *Rhizobium meliloti* bacteroids and Lucerne nodules as affected by nitrate supply. *J. Plant Physiol.* 139, 373-378.
- Aulakh, M.S., Doran, J.W., Walters, D.T., Mosier, A.R., Francis, D.D., 1991. Crop residue type and placement effect on denitrification and mineralization. *Soil Sci. Soc. Am. J.* 55, 1020-1025.
- Bohlool, B.B., Ladha, J.K., Garrity, D.P., George, T., 1992. Biological nitrogen fixation for sustainable agriculture: A perspective. *Plant and Soil* 141, 1-11.
- Hardy, R.W.F., Holsten, R.D., Jackson, E.K., Burns, R.C., 1968. The acetylene-ethylene assay for N<sub>2</sub> fixation: laboratory and field evaluation. *Plant Physiol.* 43, 1185-1207.
- IPCC, 2001. *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Kilian, S., Werner, D., 1996. Enhanced denitrification in plots of N<sub>2</sub>-fixing faba beans compared to plots of non-fixing legume and non-legumes. *Biol. Fertil. Soils* 21, 77-83.
- Klemetsson, L., Simkins, S., Svensson, B.H., Johnsson, H., Rosswall, T., 1991. Soil denitrification in three cropping systems characterized by differences in nitrogen and carbon supply. II. Water and NO<sub>3</sub> effects on the denitrification process. *Plant and Soil* 138, 273-286.
- Peoples, M.K., Craswell, E.T., 1992. Biological nitrogen fixation: Investments, expectations and actual contributions to agriculture. *Plant and Soil* 141, 13-39.
- Svensson, B.H., Klemetsson, L., Simkins, S., Paustin, K., Rosswall, T., 1991. Soil denitrification in three cropping systems characterized by differences in nitrogen and carbon supply. I. rate-distribution frequencies, comparison between systems and seasonal N losses. *Plant and Soil* 138, 257-271.
- Topp, G.C., Watt, M., Hayhoe, H.N., 1996. Point specific measurement and monitoring of soil water content with an emphasis on TDR. *Can. J. Soil Sci.* 76, 307-316.
- Zhong, Z., Hynes, R. K., Nelson, L. M., 2003. Denitrification in rhizobial species. BGSS workshop, May 29-30, Ottawa.