



Modes of plant-growth promotion of H₂-oxidizing rhizobacteria

Abdellatif, L.¹, Hamel, C.^{1,2}, Gan, Y.¹, Hanson, K.¹, Yang, C.² and Vera, C.³

¹Semi-arid Prairie Agricultural Research Centre, Swift Current, SK, Canada S9H 3X2;

²Department of Soil Science, University of Saskatchewan, Saskatoon, SK, Canada S7N 5A8;

³Melfort Agricultural Research Farm, Melfort, SK, Canada S0E1A0.



INTRODUCTION

Rhizobia trigger the formation of N₂-fixing nodules on the roots of legumes. H₂ gas is an energy-rich by-product of symbiotic nitrogen fixation. Dong and Layzell² first reported that the H₂ liberated from root nodules can promote the growth of microorganisms in the root zone, as well as plant growth. The study provides information on the plant growth promoting effect of a collection of 10 H₂-oxidizing bacteria isolated from the rhizosphere of lentil growing in semi-arid Saskatchewan. We tested the hypothesis that these lentil rhizosphere-competent bacteria can promote plant growth through various mechanisms.

MATERIALS AND METHODS

1) Characterization of plant growth promoting (PGP) trait: We tested 10 H₂-oxidizing bacteria isolated from lentil rhizosphere growing in semi-arid Saskatchewan, for phosphate-solubilizing capacity, siderophore production activity, indole acetic acid production, and ACC deaminase activity. The positive controls for the assays were from Dr. Louise Nelson, UBC, Kelowna, BC.

Screening for phosphate-solubilization capacity: The bacteria were inoculated onto K₂HPO₄-amended glucose-yeast (GY) agar medium, with an insoluble, opaque layer of calcium phosphate (Popavath et al. 2008). The experiment had a completely randomized design with six repetitions. Strain 2-106 was a positive control for comparison. Six replicates of all treatments were incubated for 5 days at 24°C in a completely randomized design (CRD). A zone of clearing around the bacterial colony indicated a positive result.

Screening for siderophore production: The bacteria were tested using the universal siderophore assay with Chrome Azurol S (CAS) and hexadecyltrimethyl ammonium bromide as indicators (Schwyn and Neilands 1987). Strain 2-106 was a positive control for comparison. Six replicates of all treatments were incubated for 5 days at 24°C in a CRD. Siderophore production was indicated by the medium color changing from blue to orange.

Screening for indole acetic acid (IAA) production: The bacteria were assayed based on the method of Patten and Glick (2002). The bacteria were inoculated on LB agar medium. Each inoculated plate was overlaid with sterile Whatman no. 1 filter paper. Strain 5-51 was a positive control. Six replicates of all treatments were incubated for 3 days at 24°C in a completely randomized design (CRD). After incubation, the filter paper was removed, and soaked in Salkowski's reagent at room temperature. A red halo indicated a positive result.

Screening for ACC deaminase activity: The bacteria were inoculated onto DF minimal salt medium (Dworkin and Foster, 1958) containing ACC as sole nitrogen source. Strain 6-8 was a positive control. Ten replicates of all treatments were incubated for 2 days at 24°C in a CRD. Growth in this medium indicates ACC deaminase activity (Husen et al. 2011).

2) Assay for growth promotion

Greenhouse assay on lentil: An experiment was conducted in growth pouches, in the greenhouse, to test the effect of the bacteria on root nodule formation. Ten bacteria plus a negative control were applied on red lentil cultivar CDC Maxim in a CRD with six replicates. The protocol of Maimaiti and Valentine (2007) was used with some modification. After 3 weeks of growth, the number of nodules per plant was recorded.

Statistical analysis: The treatment effects were statistically analyzed by ANOVA in JMP 6, using Tukey's HSD test for means comparisons.




Fig. 1 Phosphate solubilizing bacteria cultured in petri dish. The zone of clearance is seen around the colonies with the strain L1 (A). No clear zone is seen around the colonies of the strain L11 (B).

Fig. 2 Orange halos around the colonies of the strain L11 (C) and strain L22 (D) indicate the ability of those strains to produce siderophores.

Table 1 Level of different modes of action of H₂-oxidizing rhizobacterial isolates: phosphate solubilization, siderophore production, IAA and ACC deaminase activity.

Code Isolate	Isolate identity	P Solubilize (mm ²)	Siderophore (mm ²)	IAA	ACC
Control	Strain 2-106 Strain 5-51 Strain 6-8	10.3b	34.15a	+	+
L1	<i>Variovorax</i> sp.	13.9a	35.43a	+	-
L4	<i>Variovorax</i> sp.	-	25.73b	-	-
L7	<i>Rhodococcus</i> sp.	-	26.96b	-	-
L10	<i>Mycobacterium</i> sp.	-	23.55b	-	+
L11	<i>Mycobacterium</i> sp.	-	33.83a	+	+
L14	<i>Variovorax</i> sp.	-	-	-	-
L17	<i>Variovorax</i> sp.	-	26.48b	+	+
L20	<i>Acinetobacter</i> sp.	-	32.95a	-	+
L21	<i>Acinetobacter</i> sp.	-	32.53a	+	+
L22	<i>Curtobacterium</i> sp.	-	33.53a	-	+

Symbol (+): positive activity for ACC deaminase and IAA, (-) no activity for ACC deaminase and IAA.

A zone of clearing around the bacterial colony indicated phosphate solubilization. Only one bacterial strain was able to solubilize calcium phosphate (Table 1 and Fig. 1).

Nine bacterial strains produced siderophore, on the basis of change in colour of the CAS medium from blue to orange (Table 1 and Fig. 2).

Four bacterial strains were able to synthesize IAA from tryptophan. They form a red halo on the paper filter surrounding the colony in Salkowski's reagent (Table 1).

Six bacterial strains also showed positive activity for ACC deaminase when they were able to growth on DF minimal medium containing ACC (Table 1).



ACKNOWLEDGEMENTS

The authors would like to thank Heather Robinson and Fayruza Lalany for technical assistance. This research was supported by Saskatchewan Agriculture Development Fund and Saskatchewan Pulse Growers.



RESULTS




Fig. 4 Effect of the ten H₂-oxidizing rhizobacterial isolates on the number of root nodules per lentil plant three weeks after inoculation assay in the greenhouse. (A) Bars with the different letters are significantly different at $\alpha = 0.05$ (Tukey HSD, $n = 6$). (B) Lentil was inoculated with *Rhizobium leguminosarum* (Rh), and nodulation of the roots of lentil inoculated with (Rh) plus H₂-oxidizing rhizobacterial isolates. (C) Lentil inoculated with *Rhizobium leguminosarum* (Rh), and nodulation of the roots of lentil inoculated with (Rh) plus H₂-oxidizing rhizobacterial isolates (D) L14, (E) L20, and (F) L11.

➤ According to the greenhouse assays, seven H₂-oxidizing rhizobacterial have potential to promote shoot-root biomass and root nodules number per plant (Fig. 4)

CONCLUSIONS

The bacterial strains L1, L10, L11, L17, L20, L21 and L22 may have the characteristics to promote plant growth through different mechanisms (P-Solubilize, Siderophore production, IAA and ACC activity)

Greenhouse trials showed that many H₂-oxidizing rhizobacterial isolates improved the growth of lentil plants, by stimulating root nodulation. The strains L11, L17 and L20 best promoted root nodulation.

The effect of some select bacteria to promote plant growth is mainly due to phytohormones and enzymes production.

REFERENCES

- Dong Z Wu, L., Kettlewell, B., Caldwell, D., Layzell D, B. (2003) Hydrogen fertilization of soils - is this a benefit of legumes in rotation? *Plant, Cell and Environment*, 26: 1875-1879.
- Dworkin, M. and Foster, J.W. (1958) Experiments with some microorganisms which utilize ethane and hydrogen. *Journal of Bacteriology* 75, 592-603.
- Maimaiti, J., Zhang, Y., Yang, J., Cen, Y. P., Layzell, D. B., Peoples, M. and Dong, Z. (2007) Isolation and characterization of hydrogen-oxidizing bacteria induced following exposure of soil to hydrogen gas and their impact on plant growth. *Environmental Microbiology* 9(2):435-444.
- Patten, C.L. and Glick, B.R. (2002) Role of *Pseudomonas putida* Indole acetic Acid in Development of the Host Plant Root System. *Applied and Environmental Microbiology* 68, 3795-3801.
- Popavath, R.N., Gurusamy, R., Kannan Badri, N. and Sakthivel, a.N. (2008) Assessment of genetic and functional diversity of phosphate solubilizing fluorescent pseudomonads isolated from rhizospheric soil. *BMC Microbiology* 8.
- Schwyn, B. and Neilands, J.B. (1987) Universal chemical assay for the detection and determination of siderophores. *Analytical Biochemistry* 160, 47-56.
- Valentine, A.J., Osborne, B.A. and Mitchell, D.T. (2001) Interactions between phosphorus supply and total nutrient availability on mycorrhizal colonization, growth and photosynthesis of cucumber. *Scientia Horticulturae* 88, 177-189.